

Treatment of Substance Use Disorders in the Real World

Jessica M. Peirce, Ph.D.

Johns Hopkins University School of Medicine

Objectives

- identify the core components of the most common substance use disorder treatment modalities
- compare substance use disorder treatment options to make realistic and informed recommendations for patients/clients
- describe common addiction-related behaviors that patients/clients display and how to address them



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Take home points

1. You are already treating people with substance use disorders
2. You know more than you think you do



AMERICAN PSYCHOLOGICAL ASSOCIATION


9/8/2017

Definitions

- Drug = all licit and illicit drugs, alcohol; NOT tobacco
- Patients vs. clients



Figure 49. Past Year Substance Use Disorder among Adults Aged 18 or Older with Any Mental Illness in the Past Year, by Age Group: Percentages, 2008-2014



+ Difference between this estimate and the 2014 estimate is statistically significant at the .05 level.



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Figure 15. Receipt of Specialty Treatment in the Past Year among People Aged 12 or Older Who Needed Substance Use Treatment: 2014





Figure 21. Perceived Need for Substance Use Treatment among People Aged 12 or Older Who Needed Substance Use Treatment But Did Not Receive Substance Use Treatment in the Past Year: 2014



History of SUD treatment

- Developed parallel to other psychiatric treatment
- Group-based and paraprofessionals
 - *Standards for training vary widely*
- Evidence-based treatment is not uniformly offered
- Conceptualization and treatment for SUD may not reflect the highest current standards



Treatment modalities

- Adjunctive Care: NOT treatment
- Inpatient treatment
 - Short-term; Long-term
- Psychosocial outpatient & intensive outpatient
- Medication-assisted treatment
 - Buprenorphine; methadone
 - Disulfiram (Antabuse); naltrexone (Vivitrol)



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Adjunctive care

- Detoxification
- ~3 days in hospital
- Purpose is to medically manage withdrawal
- *Benzodiazepines and alcohol most common*



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Adjunctive care

- 12-step (and other) fellowships: AA, NA, Smart Recovery
- Typically regular meeting times
- Format varies, but no leader and no expectation of specific training for helpers
- Purpose is to offer social support for recovery
- *Sponsor, home group*



Adjunctive care

- Recovery housing
- Typically a group home for people with SUD
- Most are privately managed, but some are affiliated with SUD treatment
- Minimal training or regulations
- Purpose is to offer a drug-free living environment



LUXURY REHAB CENTER IN ORLANDO



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Short-term inpatient treatment

Length	7-30 days
Indication	Any SUD
Purpose	Controlled environment Establish motivation
Advantages	Removes triggers and daily stressors
Disadvantages	Can be expensive Difficult to accommodate Doesn't allow skills practice
Best for	Beginning of extended treatment



Long-term inpatient treatment


Length	3-6 months
Indication	Any SUD
Purpose	Controlled environment Establish new habits and skills
Advantages	Removes triggers and daily stressors
Disadvantages	Very few available Expensive Difficult to accommodate Doesn't allow skills practice
Best for	Beginning of extended treatment



Psychosocial outpatient/ Intensive outpatient

Length	Varies (1-3 months typical)
Indication	Any SUD
Purpose	Learn and practice skills
Advantages	Allows skills practice
Disadvantages	High dropout Quality varies widely Medication not usually offered Drug screens not universal
Best for	Extended treatment






AMERICAN PSYCHOLOGICAL ASSOCIATION

N Engl J Med 2015; 373:1789-1790

9/8/2017

17

Drug-Induced Deaths Second Only to Motor Vehicle Fatalities, 1999–2007



Source: National Center for Health Statistics, Centers for Disease Control and Prevention. National Vital Statistics Reports *Deaths: Final Data for the years 1999 to 2007 (2001 to 2010)*.



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017


18

Medication-assisted treatment

Length	Long-term (> 1 year)
Indication	Opioid use disorder (+- other SUD)
Purpose	Reduce/eliminate opioid use
Advantages	Drug screens common Highly effective
Disadvantages	May not address related problems Very long term Requires more frequent visits
Best for	Extended treatment




Medication-assisted treatment



	Buprenorphine (Suboxone, Zubsolv)	Methadone
Setting	Office or clinic	Clinic
Visits	Weekly to quarterly	Daily to 2x/mo
Counseling	Minimal	Varies
Subjective effect	Less acute effect Less withdrawal	More acute effect* Greater withdrawal
Overdose potential	Less risk	More risk

Other pharmacotherapy

- Opioids
- naltrexone (Vivitrol, Revia)
- Alcohol
- disulfiram (Antabuse): careful monitoring
- acamprosate (Campral)
- naltrexone



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

21

Evidence-based treatment: Strong

Cognitive behavioral therapy

Relapse Prevention: Marlatt; Witkiewitz

Coping with Craving: Carroll

Contingency Management: Higgins; Stitzer;
Petry

Community Reinforcement (CRA/CRAFT):
Meyers; Azrin

Behavioral Couples Therapy: O'Farrell



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Evidence-based treatment: Moderate

12-step Facilitation Therapy: Baker

Dialectical Behavior Therapy: Linehan

Acceptance and Commitment Therapy:
Hayes



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

23

Evidence-based treatment components

- Skills training and practice
- Objective and subjective monitoring of symptoms (e.g., drug use)
- Strengthen support for recovery
- Monitor treatment adherence, including medication



Elements of evidence based treatment

Joseph video #14



AMERICAN PSYCHOLOGICAL

9/8/2017

25

Treatment elements **not** recommended

- Confrontation and punishment
- Focus on “graduation”
- Reliance on self-report alone
- Discharge from treatment for drug use



But...what should I do?



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

27

What you can do

- Be aware of your own assumptions
- Conceptualize the problem accurately
- SBIRT
- Encourage harm reduction
- Use CBT principles/skills
- Consider and monitor medication
- Encourage recovery-oriented social support
- Recognize and address problems



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Stigma

the **shame** or disgrace attached to something regarded as socially unacceptable

a mark of disgrace associated with a **particular circumstance**, quality, or person



a set of negative and often unfair beliefs that a **society** or group of people have about something

Remember: WE are members of society!



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017





Life with addiction

Joseph video #13



AMERICAN PSYCHOLOGICAL

9/8/2017

31

Stigma

MORAL FAILING

Drug users are weak, lazy, sinful, immoral

Drug users are inherently flawed or broken

Drug users should be ashamed

Drug users have an addictive personality

Only the “right” treatment is appropriate

TREATMENT



AMERICAN PSYCHOLOGICAL ASSOCIATION

Stigma

COMPLEX DISORDER

Drug users have a psychiatric disorder

Some people have a chronic disorder

Drug users may do bad things, but are not bad people

Treatment should be targeted to the patient and patient needs

TREATMENT

Psychosocial and medical intervention

Need for repeated/longer treatments and continued monitoring

Take responsibility for actions, not disorder

Different or combined treatments may be necessary



AMERICAN PSYCHOLOGICAL ASSOCIATION

Conceptualization

Mood
disorder



Substance use
disorder



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Conceptualization

Mood disorder	Substance use disorder
Hiding evidence of self-harm	Lying about/hiding drug use and consequences
Refusal to participate in treatment/take medications	Refusal to participate in treatment/take medications
Belief that manic symptoms are helpful and should not be stopped	Belief that drug use is harmless or beneficial



Conceptualization

- As severity increases, treatment intensity increases
- Relapse is possible and more likely as severity increases
- Relapse is not failure of patient or treatment
- More severe and chronic disorders may need very long-term monitoring and treatment
- Medication is a tool to be used – neither the sum total of treatment nor to be discarded without consideration



SBIRT

- Screening: include a screening measure in your assessment—for *everyone*
- Brief Intervention: review your assessment, commenting on drug use
- Place drug use in context, like any other behavior
- Could end here if problem is mild



Referral to treatment

Things to consider:

- Type, frequency, amount of drugs used
- Need for medical withdrawal
- Motivation
- Social support for recovery
- Barriers to participation
- Comorbid problems



Sue, a 43 year old African-American woman, seeks help with anxious and depressive symptoms she relates to her troubled relationship. She and her husband argue several times a week; sometimes the argument gets physical. The arguments center on Sue's husband's complaints about finances and her behavior, including her drinking. Sue believes she is a social drinker, because "everyone" drinks like she does. They both drink several days a week, but more on Friday and Saturday nights when they go out with friends. Those nights, Sue has 4-5 mixed drinks. Weeknights, she will have as much as a bottle of wine.



Sue, a 43 year old African-American woman, seeks help with anxious and depressive symptoms she relates to her **troubled relationship**. She and her husband argue several times a week; sometimes the **argument gets physical**. The arguments center on Sue's husband's complaints about finances and her behavior, including her drinking. Sue **believes she is a social drinker**, because "everyone" drinks like she does. They **both drink** several days a week, but more on Friday and Saturday nights when they go out with friends. Those nights, Sue has **4-5 mixed drinks**. Weeknights, she will have as much as **a bottle of wine**.



Referral to treatment: Sue

Things to consider:

- Type, frequency, amount of drugs used
- Need for medical withdrawal
- Motivation
- Social support for recovery
- Barriers to participation
- Comorbid problems



John, a 29 year old Caucasian man, seeks help for depressive symptoms related to losing his job. He had a work-related back injury ~9 months ago and hasn't been able to work since then. John has increased his daily Oxycontin dose, but he's running out before the end of the month and the doctor is threatening to discharge him. He spends his days watching TV, smoking pot, and sleeping. John thinks he might be taking too many pills, but he also reports excruciating pain that "no one cares about." His girlfriend is fed up with his complaining and not helping around the house.



John, a 29 year old Caucasian man, seeks help for depressive symptoms related to losing his job. He had a work-related back injury ~9 months ago and hasn't been able to work since then. John has increased his daily **Oxycontin** dose, but he's **running out** before the end of the month and the doctor is threatening to discharge him. He spends his days watching TV, **smoking pot**, and sleeping. John thinks he might be taking **too many pills**, but he also reports **excruciating pain** that "no one cares about." His **girlfriend is fed up** with his complaining and not helping around the house.



Referral to treatment: John

Things to consider:

- Type, frequency, amount of drugs used
- Need for medical withdrawal
- Motivation
- Social support for recovery
- Barriers to participation
- Comorbid problems



Referral to treatment

Check SAMHSA treatment locator

Look into your local treatment centers

Ask to visit

Talk to providers

Ask about referral requirements/availability

Ask patients about their experiences*



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Find Facility

West Virginia, USA

☐ State ☐ County ☐ Distance 5 miles | [Options](#)

Service: ☒ Substance Abuse (SA) ☐ Mental Health (MH) ☐ SA & MH
☐ Health Care Centers ☐ Veterans Affairs
☐ Buprenorphine Physicians

Home


About

Services Selected

No service selected. Please select from the right.

Map

Satellite



Legend - Facility Type

- Substance Abuse
- Mental Health
- Health Care Centers
- Buprenorphine Physicians

Substance Abuse Services

Primary Focus of the Provider

[All](#) / [None](#) / [Reverse](#)

- ☐ Substance Abuse Treatment Services
- ☐ Mental Health Treatment Services
- ☐ Mix of Mental Health and Substance Abuse
- ☐ General health services

Type of Care

[All](#) / [None](#) / [Reverse](#)

- ☐ Substance abuse treatment
- ☐ Detoxification
- ☐ Methadone maintenance
- ☐ Methadone maintenance for predetermined time
- ☐ Methadone detoxification
- ☐ Buprenorphine maintenance
- ☐ Buprenorphine maintenance for predetermined time
- ☐ Buprenorphine detoxification
- ☐ Relapse prevention from naltrxone
- ☐ Buprenorphine used in treatment
- ☐ Naltrexone (oral)
- ☐ Vivitrol® (injectable Naltrexone)
- ☐ Methadone




AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

YOUR TITLE HERE

46



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

47

Availability

	Methadone providers	Buprenorphine providers*
West Virginia	9	250
New Hampshire	8	123
Kentucky	12	500

* 2/3 or more of physicians with a buprenorphine waiver do not write any prescriptions



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Within your treatment

- Harm reduction
- Reducing use
- Safety measures: Narcan, informing others of use
- CBT principles/skills training
- Motivational approaches (MI/MET)
- Triggers: “people, places, and things”
- Coping strategies



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Within your treatment

- Medications
 - Addiction medications
 - Consider SUD when thinking about other medications
 - State PDMP
- Social support for recovery
 - Establish contact with family/other treatment providers
 - Social skills and practice



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

50

The role of supports in recovery

Joseph video #19



AMERICAN PSYCHOLOGICAL

9/8/2017

51

But...what if *they*...?



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

52

Problem situations

Active drug use interfering with recovery goals

- Denial of drug use

- Acute drug use

- Ongoing drug use

Nonadherence to treatment goals

- Refusal to attend treatment

- Inadequate treatment

Concern about relapse



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Active drug use

Acute vs. chronic drug use

Consider alcohol, prescription and illicit drugs

Use non-stigmatizing language

“Drug-affected” instead of “high”

Describe concerning objective behaviors

“Your eyes are closing while I’m talking, and I’m worried you’re not able to fully focus on this meeting.”

“You haven’t attended the medical appointments that you said you would.”



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Active drug use

Ask!

“What do you think the problem is?”

“The benzodiazepines you’re taking could be affecting your memory. How many have you taken?”

“Have you taken anything that could be affecting you in that way?”

Join together to solve the problem

“If taking those medications makes you so tired, what do you think you could do differently?”



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Active drug use

Don't engage with severely drug-affected people

Be very directive; focus on safety

Postpone discussion about concerns until they are alert and aware

Potential for overdose if

Not arousable

Acute medical problem: vomiting, seizures

Get immediate help

Acute drug use is similar to acute suicidality!



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Nonadherence to treatment

Motivational approach to explore ideas

Encourage SUD treatment if needed

Support SUD treatment goals

- Attendance for counseling, medication

- Self-help groups

Offer to coordinate care

- Special release of information



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Concern about relapse

Lapses and relapses are common and don't indicate failure

Ask!

Cravings, triggers, any lapses

Provide support

Engagement/reengagement with treatment

Shorter is better – help to shorten relapses!



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

What you can do

- Be aware of your own assumptions
- Conceptualize the problem accurately
- SBIRT
- Encourage harm reduction
- Use CBT principles/skills
- Consider and monitor medication
- Encourage recovery-oriented social support
- Recognize and address problems



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

Take home points

1. You are already treating people with substance use disorders
2. You know more than you think you do



AMERICAN PSYCHOLOGICAL ASSOCIATION

9/8/2017

60

Thank you

Jessica M. Peirce, Ph.D.



AMERICAN PSYCHOLOGICAL

9/8/2017

61