

Disordered Gaming: The Role of a Gamer's Distress Profile

Jade Kovacs¹, Daniel Zarate², Gabriel de Sena Collier¹, Thong Thai Diep Tran¹, and
Vasileios Stavropoulos^{2, 3}

¹ College of Health and Biomedicine, Victoria University

² Institute for Health and Sport, Victoria University

³ Department of Psychology, University of Athens

Internet Gaming Disorder (IGD) embodies a persistent and recurrent engagement with video games, to the exclusion of other activities, that cannot be controlled, and with significant impairments in everyday functioning. Previous research suggests that IGD is experienced differently depending on the gamer's profile, while distress symptoms such as depression, anxiety, and stress have been independently associated to the development of IGD. Interestingly, no study to date has aimed to profile gamers based on these three psychopathologies. The present study aimed to (a) profile gamers concerning their depression, anxiety, and stress, and (b) examine the differences in IGD levels between the different profiles of distress. A sample consisting of 968 gamers (18–64 years, $M_{age} = 29.54$) was assessed with the Depression, Anxiety and Stress Scale (DASS-21) and the Internet Gaming Disorder Scale–Short-Form (IGDS9-SF). Latent class analysis (LCA) identified three distinct distress profiles. These encompassed “High-Distress Comorbidity” (HDC; 25.9%), “Medium-Distress Comorbidity” (MDC; 48.7%) and “Low-Distress Comorbidity” (LDC; 25.4%) gamers. As hypothesised, higher distress comorbidity profiles are linked with higher IGD levels. Findings suggest that there are different distress profiles varying by symptom severity, with more distressed gamers presenting with higher IGD levels independent of the specific distress symptom they reported. Thus, differential diagnosis of depression, anxiety, and stress symptoms may not be necessary in the context of IGD treatment, but rather distress behaviours should be targeted concurrently and independent of their nature.

Public Significance Statement

Both the type of gamer a person is and their level of distress play a part in the likelihood of developing Internet Gaming Disorder (IGD). Different profiles of distress were examined in relation to IGD levels, with results showing that higher distress profiles were accompanied by higher IGD levels.

Keywords: anxiety, depression, Internet Gaming Disorder, latent class analysis, stress

Internet gaming has attracted research interest due to its prevalence among all age-groups and its potential effects on psychological health and well-being (Anderson et al., 2017; Stavropoulos, Anderson, et al., 2018; Stavropoulos, Beard, et al., 2018). Previous research has shown that gaming can facilitate a wide range of cognitive, therapeutic, and

social benefits (Granic et al., 2014; Raith et al., 2021). However, excessive gaming has been associated with anxiety, depression, obsessive-compulsive behaviour, attention deficit/hyperactivity, social phobias, and loneliness (Männikkö et al., 2020; Tullett-Prado et al., 2021; Wong et al., 2020). These negative repercussions

This article was published Online First May 5, 2022.

Vasileios Stavropoulos  <https://orcid.org/0000-0001-6964-4662>

Vasileios Stavropoulos received funding from the Victoria University Early Career Researcher Fund ECR 2020, Number 68761601, and the Australian Research Council, Discovery Early Career Researcher Award, 2021, Number DE210101107. The authors of the present study do not report any conflict of interest.

All procedures performed in the study involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. This article does not contain any studies with animals performed by any of the authors. The authors would thank Daniel Zarate for his unconditional support and guidance during this adventurous learning process.

Jade Kovacs authored the main body, contributed to the literature review, data collection, framework formulation, data analysis and the

structure and sequence of theoretical arguments. Daniel Zarate contributed to the literature review, framework formulation, data analysis and the structure and sequence of theoretical arguments. Gabriel de Sena Collier contributed to data collection, data analysis and proofread the final manuscript. Thong Thai Diep Tran contributed to data collection, formatting and proofread the final manuscript. Vasileios Stavropoulos contributed to the theoretical consolidation of the present work and revised and edited the final manuscript.

The data and syntax that support the findings of this study are available on the following two github links: Data are available at <https://github.com/Vas08011980/Distress-Profiles-IGD/commit/49862759f96c34e632a0c137328cd44e7a30e476>. Syntax is available at <https://github.com/Vas08011980/Distress-Profiles-IGD/tree/main>.

Correspondence concerning this article should be addressed to Vasileios Stavropoulos, Institute for Health and Sport, Victoria University, P.O. Box 14428, Melbourne 8001, Australia. Email: vasileios.stavropoulos@vu.edu.au

have prompted scholars to introduce diagnostic classifications related to disordered gaming (Stavropoulos et al., 2020). For example, in 2013, the American Psychiatric Association included Internet Gaming Disorder (IGD) in the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5) under Section III as a condition warranting more clinical research and experience before it may be considered as a formal disorder in the primary manual (DSM-5; American Psychiatric Association, 2013). Five years later (2019), Gaming Disorder (GD) was recognized as an official classification in the 11th edition of the *International Classification of Diseases* (ICD-11) by the World Health Organization (World Health Organization, 2019).

According to the DSM-5, IGD consists of nine core symptoms with at least five of these present over 12-months (American Psychiatric Association, 2013). These nine clinical symptoms are (a) a preoccupation with video games (preoccupation); (b) withdrawal symptoms (e.g., irritability and frustration) in the absence of gaming (withdrawal); (c) the need to spend an increased amount of time playing video games to create the same “high” (tolerance); (d) an inability to control how much time is spent playing video games, even when the subject wants and attempts to (loss of control); (e) a loss of interest in previously enjoyed activities (other than gaming) as a consequence of video games (surrendering from other activities); (f) continuing to play video games despite understanding their negative effects (continuation); (g) deceiving those close to them, therapists or others regarding their gaming habits (fraud); (h) utilizing video games to deal with or eliminate negative feelings (escape), and finally; (i) damage or loss of relationships, work or education as a result of video game engagement (negative consequences-reducing one's everyday life functionality).

Conceptualization of Disordered Gaming

Individual differences interwoven with different risk profiles have been suggested to contribute to the development of IGD. Particularly, distress symptoms such as depression and anxiety (along with other factors) may act as risk factors for IGD (Burleigh et al., 2018; Liew et al., 2018; Stavropoulos, Motti-Stefanidi, et al., 2021). There have been many explanatory proposals put forward to understand how such individual differences may interact with each other in relation to IGD (Brand et al., 2016; Davis, 2001; Griffiths, 2005; Griffiths et al., 2017; Kardefelt-Winther, 2014). For example, the self-medication hypothesis (Khantzian, 1997) has been adopted to understand behavioural addictions including IGD and posits that distressed individuals may engage in problematic online gaming as a means of relief (Griffiths, 2005; Griffiths et al., 2017). Similarly, the compensatory internet use (CIU) model (Kardefelt-Winther, 2014) suggests that disordered gaming may be a coping strategy to manage psychopathological symptoms or negative life events (although concerns regarding the over pathologizing of common behaviours have been raised; Kardefelt-Winther et al., 2017; van Rooij et al., 2018). Much like the self-medication hypothesis, this model posits the idea that addictive use of the internet may help one cope with their psychological suffering such as anxiety, depression, and stress (Griffiths, 2005; Griffiths et al., 2017).

While aetiological models (e.g., CIU) provide useful potential explanations, many researchers have suggested an alternative “typological” approach to understanding the patterns in which IGD may manifest (Billieux, Schimmenti, et al., 2015; Billieux,

Thorens, et al., 2015; Colder Carras & Kardefelt-Winther, 2018; Faulkner et al., 2015; Lee et al., 2017; Lemmens et al., 2015; Pontes et al., 2014; Tullett-Prado et al., 2021; Ünlübol et al., 2020). This approach focuses on identifying variability across different profiles of gamers based on IGD symptoms or behaviours to better understand the gamers' context (Stavropoulos et al., 2020; Stavropoulos, Gomez, et al., 2021; Stavropoulos, Motti-Stefanidi, et al., 2021; Stavropoulos, Vassallo, et al., 2021).

Such variability among gamers' profiles has been observed via latent profile analyses (LPA) considering IGD-related symptomatology (Billieux, Schimmenti, et al., 2015; Billieux, Thorens, et al., 2015; Colder Carras & Kardefelt-Winther, 2018; Faulkner et al., 2015; Lee et al., 2017; Lemmens et al., 2015; Pontes et al., 2014; Tullett-Prado et al., 2021; Ünlübol et al., 2020). For example, Pontes et al. (2014) used the 20-item IGD and identified five distinct profiles in a population of gamers ($n = 1,003$) where profiles varied according to symptom severity and gameplay hours (i.e., casual, regular, low-risk, high-risk, and disordered gamers). Similarly, Colder Carras and Kardefelt-Winther (2018) used the Assessment of Internet and Computer game Addiction–Gaming Module on a population of 7,865 adolescent gamers from Europe and identified five varying profiles of gamers (i.e., normative, IGD, concerned, at risk, engaged). Interestingly, both anxiety and depression significantly predicted membership of the IGD “engaged” and “concerned” profiles. Lemmens et al. (2015) utilized the Internet Gaming Disorder Scale-Short-Form and identified three profiles (e.g., normal, risky, disordered) based on time spent gaming, self-esteem, loneliness, and aggression. Additionally, Faulkner et al. (2015) utilized the Problem Video Gameplaying Scale on a population of 3,338 high-school students in the USA aged between 11 and 20 years. Informed by the severity of problematic gaming, they identified four profiles (e.g., normative, low, high, severe). Students in the severe profile had significantly more depression and anxiety symptoms than students in the high, low, and normative profiles. While these profiles have been observed considering IGD-related symptomatology, they have not been described/portrayed, to the best of the author's knowledge, on the basis of one's comorbid distress symptoms as potential risk for the development of IGD.

Distress and Disordered Gaming

Consistent with the aforementioned self-medication (Khantzian, 1997) and CIU (Kardefelt-Winther, 2014) models, “negative escapism” suggests gaming is negatively reinforced as a means of avoiding symptoms of distress (Martín-Fernández et al., 2016). Indeed, individuals with depression, anxiety, or stress may spend excessive time gaming on the internet as a coping mechanism to deal with worries and difficulties in their life (Ho et al., 2014; Yen et al., 2019). Similarly, IGD and low mood may often present with several behaviours in common, including social withdrawal, fatigue, disruption of sleep, and poor performances in school and work (Achab et al., 2011).

Subsequently, in terms of IGD comorbidities, depression, anxiety, and stress are among the most frequently reported health-related variables associated with IGD (Darvesh et al., 2020; Wong et al., 2020). A recent meta-analysis investigating depression in individuals diagnosed with IGD observed that the comorbidity of depression in individuals with IGD ranged up to 75% with an average prevalence of 32% (Ostinelli et al., 2021). Similarly, a meta-analysis examining the relationship between IGD and comorbid psychopathologies

found that 92% (out of 13) of studies involving anxiety, described significant positive correlations between anxiety and IGD (González-Bueso et al., 2018). Furthermore, stress has also been shown to have a strong association with IGD suggesting that stress is a risk factor for addiction and increases the likelihood of relapses (Goeders, 2003; González-Bueso et al., 2018; Stavropoulos, Vassallo, et al., 2021).

Present Study

The above-described research indicates a relationship between psychological distress and IGD; however, the details of this relationship are not yet entirely elucidated. In particular, the degree to which the individual facets of psychological distress are seen in individuals of varying IGD experience has yet to be examined. Given the connection between distress and IGD, and the heterogeneity of IGD experience, this makes a worthwhile avenue of investigation into the risks and consequences of IGD. This is particularly beneficial as comorbid psychopathology may require to be treated concurrently with IGD for optimum results (Zajac et al., 2020).

Accordingly, the present study aspires to address the following. Firstly, to advance past knowledge considering the typologies/profiles of distress symptoms within the gaming population. (e.g., can an online community sample of gamers be described by different distress profiles/typologies?). Secondly, to expand the empirical evidence considering distress symptoms as potential IGD aetiology (e.g., is there a significant difference in IGD levels between the different profiles of distress?). These aims will be innovatively addressed by the examination of a large cohort of gamers ($N > 900$), the use of a psychometrically sound and broadly used distress scale (i.e., Depression, Anxiety and Stress Scale-21 items; DASS-21; Lovibond & Lovibond, 1995) and the implementation of a statistically advanced sequence of 12 potential profiling models (Rosenberg et al., 2019). Accordingly, to address the outlined aims, the following research questions were elaborated:

Research Question 1: What is the number and nature of distress profiles that best describes the cross-sectional sample examined, taking into consideration levels of depression, anxiety and stress?

Research Question 2: What is the proportion of individuals in each profile based on the selected indicators?

Additionally, the following hypothesis was elaborated:

Hypothesis 1: Participants experiencing higher levels of distress will display higher levels of disordered gaming.

Method

Participants

An online community sample of English-speaking, adult gamers aged 18–64 were included in the study. The initial sample comprised 1,097 responses, with seven participants deleted due to being preview-only responses, five deleted after being flagged as spam responses, and 11 deleted due to being potential bots, 11 deleted as they did not provide consent to participate in the study, 16 were deleted for not providing their age, two were deleted for being younger than 18, and a further 77 were excluded due to providing

minimal responses to the battery of questionnaires. After this data screening process, the sample consisted of $N = 968$, $M_{\text{age}} = 29.54$, $SD_{\text{age}} = 9.35$, Males = 622 (64.3%), Females = 315 (32.5%), other = 31 (3.2%, including trans/nonbinary, genderqueer). Missing values in participants' responses represented 0.12% and were below recommended thresholds ($>5\%$; Schafer, 1999) and missing completely at random (MCAR; Little's test $\chi^2 = 314.979$, $df = 281$, $p = .080$). Therefore, we proceeded with the analyses. Table 1 shows participants sociodemographic information.

Measures

Depression, Anxiety, and Stress Scale (DASS-21) was utilized to assess, via self-reporting, depression, anxiety, and stress with seven items for each self-report subscale (Lovibond & Lovibond, 1995). The 21 items are rated on a 4-point Likert scale ranging from 0 to 3 with a total score for each subscale ranging between 0 and 21. Examples of items include "I found it hard to wind down." Higher DASS scores indicate a higher level for that corresponding subscale. The DASS-21 has showed excellent internal reliability (Cronbach's $\alpha = 0.81$, 0.89 and 0.78; Coker et al., 2018) and appropriate convergent validity (Le et al., 2017) in previous research and the present study (Stress Cronbach $\alpha = .883$, and McDonald's $\omega = .883$, anxiety $\alpha = .865$, $\omega = .869$ depression $\alpha = .931$, and $\omega = .932$ and for the whole scale (distress factor) $\alpha = .950$, and $\omega = .950$).

Internet Gaming Disorder Scale-Short-Form (IGDS9-SF) is a short psychometric self-report tool to examine, via self-reporting, online and offline gaming activities within the last 12 months. The nine items are measured on a 5-point Likert scale ranging from 1 (*Never*) to 5 (*Very Often*) with total scores ranging from 9 to 45, and higher scores indicating higher levels of IGD. Examples of items include "Do you feel preoccupied with your gaming behaviour." The IGDS9-SF shows acceptable internal reliability with a Cronbach's α of 0.87, as well as high convergent validity (Kim & Ko, 2020). For the present study, the internal reliability rate of the scale was Cronbach $\alpha = .885$, and McDonald's $\omega = .892$.

Procedure

Upon obtaining approval from the Victoria University Human Research Ethics Committee (HRE20-169), participants were recruited through convenience sampling. A Qualtrics link was distributed through social media platforms. The survey was targeted towards gaming populations by using keywords such as internet games, PC gamer, massive multiplayer online role playing gamers (MMORPGs), multiplayer online battle arena (MOBAs), first player shooter (FPS). Upon accessing the link, participants were directed to the Plain Language Information Statement clearly stating the study aims, potential contributions, voluntary participation, right to withdraw, and informed consent. Informed consent was assured as participants were required to tick a box prior to beginning the survey. Considering that the present study aimed to protect participants from inadvertent harm, all participant information remained anonymous with individual participants only being identifiable by number.

Statistical Analyses

To address RQ₁ and RQ₂, the depression, anxiety, and stress subscales as assessed by the DASS-21 were employed as indicators

Table 1*Sociodemographic Information and Descriptive Statistics Across Gender Groups*

Demographic information	Gender		
	Female	Male	Other
Employment status			
Full-time	86	238	7
Part-time	49	61	1
Casual	11	12	0
Self-employed	17	48	2
Retired	2	3	0
Unemployed	58	122	7
Full-time student	43	92	6
Other	49	46	8
Sexual orientation			
Heterosexual/straight	211	529	3
Homosexual/gay	13	33	4
Bisexual	65	48	12
Unidentified/other	26	12	12
Ethnicity/background			
Black/African American	23	31	1
White/Caucasian	193	380	22
Asian	59	124	1
Hispanic/Latino	9	35	2
Aboriginal/Torres Strait Islander	1	0	0
Indigenous	1	1	1
Indian	1	4	0
Pacific Islander	1	3	0
Middle-Eastern	2	2	0
Mixed	25	40	3
Other	0	2	1
Romantic relationship			
Yes	187	247	17
No	118	356	14
Prefer not to say	10	19	0
Education			
Elementary or middle school	2	10	0
High school or equivalent	74	166	11
Vocational/Technical and further education (TAFE)	26	55	4
Some tertiary education	69	113	3
Bachelor's degree (3 years)	76	137	5
Honours degree or equivalent (4 years)	35	69	5
Postgraduate degree (PhD, MS, etc.)	30	59	2
Other	3	13	1
Marital status			
Single	164	405	23
Living with another	62	68	7
Married	68	120	0
Separated	2	4	0
Divorced	10	10	0
Widowed	2	1	0
Other	7	14	1
Descriptive statistics			
Depression	7.57 (5.80)	8.82 (5.98)	10.6 (6.58)
Anxiety	4.29 (4.16)	5.86 (4.79)	6.84 (5.37)
Stress	6.33 (4.66)	8.56 (4.90)	10.3 (5.58)
IGD	18.60 (7.26)	17.30 (6.72)	18.5 (7.53)

Note. IGD = Internet Gaming Disorder.

for a LPA using the TIDYLPa CRAN package in R (Rosenberg et al., 2019). This analysis was chosen for its modelling approach which allows for the identification of naturally homogenous sub-groups (profiles) within a population based on descriptors or characteristics of significance (Muthén & Muthén, 2016). Specifically, LPA employs a Maximum Likelihood Estimator (MLE) to identify profile membership probabilities among gamers based on their distress symptoms. TidyLPA was selected for its ability to estimate optimal relationships between indicators across different profiles, including means (i.e., average levels of DASS), variances (i.e., variability of DASS within profiles), and covariances (i.e., variability of DASS across profiles; Tullett-Prado et al., 2021). Table 2 below shows the four possible combinations of parameterizations of variance-covariance structures that can be estimated with TidyLPA to obtain the optimal number of profiles (for a thorough explanation see Masyn, 2013, p. 585).

Selecting the optimal number of latent profiles involved a sequential process. Firstly, identification of the best combination of parameters, including (un)constrained profile mean, variance, and covariance, by comparing models based on the Akaike Information Criterion (AIC), Bayesian Information Criterion (BIC), Approximate Weight of Evidence Criterion (AWE), Classification Likelihood Criterion (CLC), and Kullback Information Criterion (KIC) with smaller values indicating better fit (Masyn, 2013). Secondly, assessing the best number of profiles via the bootstrapped likelihood ratio test (BLRT) to determine if adding an extra latent profile resulted in a significant increase in fit (with $p < .05$ as indication of improved fit; McLachlan, 1987). Finally, evaluation of standardized entropy criterion (h) to assess heterogeneity levels across latent profiles, with 0.40–0.60 indicating low, 0.60–0.80 medium, and $> .80$ high entropy (Celeux & Soromenho, 1996; Clark & Muthén, 2009).

Additionally, to address H_1 , a one-way analysis of variance (ANOVA) was conducted to examine the difference in IGD scores between the different distress profiles. Further post hoc analyses were conducted to examine where these differences lied.

Results

Identifying and Describing Distress Profiles

To answer RQ₁ and RQ₂, we sought to identify the optimum number of latent profiles and the population share in each profile. Table 3 shows initial testing of 24 possible combinations of models, varying by number of classes and parametrization. The Class Variant Diagonal Parameterization (CVDP) model with three profiles and the Class Variant Unrestricted Parameterization (CVUP) model with two profiles were further examined due to their lower AIC and BIC values.

Table 4 shows further testing to expand fit indices for the CVDP model with three profiles and the CVUP model with two profiles (Rosenberg et al., 2019). Both options showed significant BLRT- p and an appropriate number of participants in the smallest latent profile. While the CVUP model with two profiles yielded superior AIC and BIC, the CVDP model with three profiles resulted in better level of classification accuracy (entropy = 0.84) and was therefore selected due to optimum fit. Observed entropy for this model significantly exceeded the cut-off point of 0.76 (Larose et al., 2016),

Table 2*Parameterization of Variance–Covariance Structures, From the Most to the Least Restrictive Model*

Model	Variances	Covariances	Parameterization type
1	Equal	Fixed to 0	Class-invariant diagonal parameterization model (CIDP). This model assumes that relationships across model indicators should not be estimated (covariances fixed at zero) and that different profiles will be qualitatively similar (equal variances).
2	Varying	Fixed to 0	Class-varying diagonal parameterization model (CVDP). This model assumes that relationships between model indicators should not be estimated (covariances fixed at zero), and that different profiles will be qualitatively different (varying variances).
3	Equal	Equal	Class-invariant unrestricted parameterization model (CIUP). Indicators are allowed to co-vary within profiles, and the variances and covariances are restricted to be equal across different profiles.
4	Varying	Varying	Class varying unrestricted parameterization (CVUP). All the indicators are allowed to co-vary within profiles, and the variances and covariances (i.e., residual correlations) are allowed to be different across profiles. In other words, this model assumes that there are relationships between model indicators within and between latent profiles that should be estimated (i.e., varying covariances), and that different profiles will be qualitatively different (varying variances).

Note. In this context, “diagonal” indicates that the sum of elements in the variance–covariance matrix equals to zero, thus effectively preventing the model to estimate covariances between indicators.

suggesting that the accurate classification of the CVDP 3 profile structure was over 90% correct (Larose et al., 2016). Accordingly, the share of participants in each estimated profile were $n = 251$ (25.9%) for Profile 1, $n = 471$ (48.7%) for Profile 2, and $n = 246$ (25.4%) for Profile 3. Table 5 displays the profiles standardized mean scores, raw mean scores, and standard deviations of depression, anxiety, and stress.

Distress latent profiles were described considering both raw and standardized reported symptoms to examine their distinct features while concurrently enabling objective understanding in the terms of normal distributions of depression, anxiety, and stress (RQ₁). In that context, the three latent distress profiles showed variability in raw scores and mean values of depression, anxiety, and stress levels. Figure 1 illustrates mean differences in distress symptoms across

Table 3*Initial Model Testing*

Model	Profiles	AIC	BIC	AWE	CLC	KIC
Class invariant	1	8247.963	8277.215	8334.466	8237.963	8256.963
Diagonal	2	7120.837	7169.589	7266.682	7102.496	7133.837
Parameterization (CIDP)	3	6677.503	6745.757	6882.371	6651.142	6694.503
	4	6496.618	6584.372	6462.277	6517.618	6760.468
	5	6418.474	6525.729	6741.365	6376.093	6443.474
	6	6426.426	6553.182	6808.477	6375.887	6455.426
Class variant	1	8247.963	8277.215	8334.466	8237.963	8256.963
Diagonal	2	6848.087	6911.465	7038.185	6823.744	6864.087
Parameterization (CVDP)	3	6251.775	6349.280	6545.110	6213.450	6274.775
	4	N.C	N.C	N.C	N.C	N.C
	5	N.C	N.C	N.C	N.C	N.C
	6	N.C	N.C	N.C	N.C	N.C
Class invariant	1	6636.920	6680.797	6767.674	6620.920	6648.920
Unrestricted	2	6645.865	6709.243	6837.419	6620.067	6661.865
Parameterization (CIUP)	3	6467.436	6550.315	6716.675	6434.956	6487.436
	4	6415.292	6517.672	6723.420	6374.924	6439.292
	5	6270.024	6391.904	6637.278	6221.531	6298.024
	6	6278.008	6419.390	6704.482	6221.297	6310.008
Class variant	1	6636.920	6680.797	6767.674	6620.920	6648.920
Unrestricted	2	6167.752	6260.381	6446.322	6131.440	6189.752
Parameterization (CVUP)	3	N.C	N.C	N.C	N.C	N.C
	4	N.C	N.C	N.C	N.C	N.C
	5	N.C	N.C	N.C	N.C	N.C
	6	N.C	N.C	N.C	N.C	N.C

Note. This table shows comparisons between different number of profiles for four possible combination of model parameters (including varying/fixed classes and varying/fixed covariances). Highlighted results (bold) indicate best model parameterization according to the best information criterion. Results showing N/C indicate that no convergence on a solution was possible. AIC = Akaike Information Criterion; BIC = Bayesian Information Criterion; AWE = Approximate Weight of Evidence Criterion; CLC = Classification Likelihood Criterion; KIC = Kullback Information Criterion; CIDP = Class-invariant diagonal parameterization model; CVDP = Class-varying diagonal parameterization model; CIUP = Class-invariant unrestricted parameterization model; CVUP = Class varying unrestricted parameterization.

Table 4*Fit Indices of CVDP With Three Profiles and CVUP With Two Profiles*

Model	Profiles	AIC	BIC	Entropy	Proportion of smallest profile	BLRT-p
CVDP	3	15671.03	15768.53	0.84	0.25	0.01
CVUP	2	15605.49	15698.12	0.74	0.34	0.01

Note. This table shows that CVDP model with three latent profiles demonstrate a higher entropy value resulting in better differentiation between profiles. AIC = Akaike Information Criterion; BIC = Bayesian Information Criterion; BLRT-p = Bootstrapped likelihood ration test; CVDP = Class Variant Diagonal Parameterization; CVUP = Class-Variant Unrestricted Parameterization. Bold values indicate optimum model.

latent profiles, and Figure 2 compares raw distress symptoms with normal population scores (as suggested by Coker et al., 2018). Individuals classified in *Profile 1* scored in the “extremely severe” range for depression (14+) and anxiety (10+), “severe” range for stress (13–16), and scored +1.15SD to +1.26SD with respect to mean distress values for our sample. Consequently, Profile 1 was defined as “High-Distress Comorbidity” (HDC), distinguished by the highest distress symptoms. Participants in *Profile 2* scored in the “moderate” range for depression (7–10), “mild” range for anxiety (4–5), “normal” range for stress (0–7), remained near mean distress levels (−0.17SD to −0.05SD), and were thus labelled “Medium-Distress Comorbidity” (MDC). Finally, participants in *Profile 3* scored in the “normal” range for depression (0–4), anxiety (0–3), stress (0–7), and scored below mean levels (−1.10SD to −0.92SD). This profile was thus defined as the “Low-Distress Comorbidity” profile (LDC), distinguished by the lowest symptom experiences when compared with the other distress profiles.

Hypothesis 1: Distress Profiles and IGD

With α set at .05, a one-way ANOVA without assuming homogeneity of variance (i.e., Welch’s test) was conducted to examine the differences between IGD scores within Profile 1 (HDC), Profile 2 (MDC), and Profile 3 (LDC). The results indicated that there was a significant and large effect on IGD scores between the different profiles, $F_{\text{Welch}}(2, 528.15) = 101.94, p < .001$. Games–Howell post hoc analyses revealed that the differences in IGD were significant across all three different profiles (high distress to mid distress; $p < .001$; SE: .59; CI [2.03, 4.85]; high distress to low distress; $p < .001$; SE: .59; CI [6.32, 9.14]; mid distress to low distress; $p < .001$; SE: .42; CI [3.31, 5.28]. In other words, the mean IGD scores within the profiles steadily decreased in descending order of HDC, MDC, and LDC (see Figure 3). Interestingly, those belonging to the “High-Distress Comorbidity” profile also displayed a tendency to score higher in IGD compared with individuals in other profiles (observed by elevation and size of dots). This supports our hypothesis indicating that profiles with concurrently higher depression, anxiety and stress are accompanied by higher levels of IGD.

Discussion

The present study was the first to examine distress profiles in a large sample of 968 gamers. This was done via the use of a psychometrically sound and broadly used distress scale (i.e., DASS) and the implementation of a statistically advanced sequence of 24 potential profiling models (Rosenberg et al., 2019). Findings suggested the presence of three distinct distress profiles. These encompassed “High-Distress Comorbidity” (HDC; 25.9%), “Medium-Distress Comorbidity” (MDC; 48.7%) and “Low-Distress Comorbidity” (LDC; 25.4%) gamers. As hypothesised, higher distress comorbidity profiles were associated with higher IGD levels.

Different Distress Profiles Among the Gaming Population

The findings suggested three distinct profiles of distress present within this gaming population. The depression, anxiety, and stress levels within each profile were qualitatively different. Specifically, participants in the HDC profile showed severe stress, extremely severe depression, and anxiety, and ranged between 1.15 and 1.26 SDs above mean sample scores. Participants in the MDC profile showed moderate depression, mild stress, normal anxiety, and remained within sample mean levels. Finally, participants in the LDC scored in the normal range for depression, anxiety and stress, and steadily ranged 0.9–1.10 SDs below the sample mean. This indicates that participants in the HDC profile were at higher risk of experiencing distress symptoms compared with other latent profiles and expected normal population scores.


These findings are consistent with previous research which has demonstrated gamers do not encompass a homogenous group, but rather a group consisting of multiple different profiles identifiable by unique characteristics (Billieux, Schimmenti, et al., 2015; Billieux, Thorens, et al., 2015; Colder Carras & Kardefelt-Winther, 2018; Faulkner et al., 2015; Lee et al., 2017; Lemmens et al., 2015; Pontes et al., 2014; Tullett-Prado et al., 2021; Ünübol et al., 2020). Interestingly, the presentation of such profiles suggest that most gamers (48.7%) represent a normative

Table 5*Description of Distress Profiles Including Population Share, and Raw and Standardized Mean Scores of Depression, Anxiety, and Stress*

Profile	N	%	Depression	Z depression	Anxiety	Z anxiety	Stress	Z stress
High comorbidity	251	25.9	14.90 (4.16)	1.15	10.30 (4.23)	1.22	13.40 (3.47)	1.26
Mid comorbidity	471	48.7	7.76 (3.95)	−0.05	4.13 (2.44)	−0.17	6.70 (2.39)	−0.17
Low comorbidity	246	25.4	1.77 (1.88)	−1.06	0.75 (0.91)	−0.92	1.76 (1.64)	−1.10

Note. Z scores represent standardizes scores and standard deviation is presented between brackets.

Figure 1
Distress Latent Profiles



Note. The plot illustrates three distinct latent profiles considering participants' symptoms of distress measured in standard deviation from the mean including depression, anxiety, and stress. The high line represents participants experiencing high levels of comorbid distress symptoms, the middle line medium levels, and the lower line low levels. See the online article for the color version of this figure.

distribution of distress within the gaming population. This finding adds valuable evidence in support of the de-pathologizing of the gaming population, which has often been pathologized and is an ongoing topic of debate within the literature (O'Brien, 2018; van Rooij et al., 2018). In other words, results support that being a gamer does not necessarily correspond with high/pathological levels of distress behaviours, such as depression anxiety and stress, examined in the present study.

Such findings are particularly relevant considering recent events wherein the Chinese government imposed further restrictions to minors' online video gameplay due to concerns surrounding gaming addiction (Goh, 2021). As of August 2021, the government restricted game time to a limit of 1 hr of play only on Friday, Saturday, and Sunday, in addition to increasing the frequency and intensity of government inspections surrounding online gaming companies to ensure these restrictions are followed (Goh, 2021). Indeed, such policies are congruent with the idea that gaming is inherently negative, disregarding the role of gaming in normal development and how differences between gamers may relate to their unique experiences over time (Stavropoulos, Gomez, et al., 2021).

Relationship Between Distress Profiles and IGD Levels

In line with previous research, the present study reported a significant association between one's IGD levels and their distress levels (Burleigh et al., 2018; Faulkner et al., 2015; Goeders, 2003; Lemmens et al., 2015; Liew et al., 2018; Wong et al., 2020). These findings are further supported when considering the aetiological models put forth to explain the underlying causes of IGD. For example, the self-medication (Griffiths, 2005; Griffiths et al., 2017) and CIU model (Kardefelt-Winther, 2014) propose that distressed individuals may engage in problematic gaming behaviour to cope with psychological suffering resulting in "negative escapism" (Martín-Fernández et al., 2016; Valentino et al., 2010). Indeed,

Lee et al. (2017) speculated that emotionally vulnerable individuals may engage in internet gaming for mood modification purposes that potentially stem from underlying depression or external stressors. Similarly, the cognitive-behavioural theory of pathological internet use (Davis, 2001) and the Person-Affect-Cognitive-Execution (I-PACE) model (Brand et al., 2016) suggest that depression, anxiety, or stress can be contributing factors (either precipitating and/or perpetuating) regarding the development of IGD.


Considering the above, findings identified in the present study pose significant clinical practise directions. Specifically, observed distress profiles suggest that gamers with simultaneously higher levels of depression, anxiety, and stress may have a greater likelihood of developing IGD, and thus should be considered at risk and prioritized in prevention, intervention, and primary care. This implies that differential diagnosis processes (i.e., differentially assessing/distinguishing symptoms of depression, anxiety, stress, and disordered gaming) should be followed when assessing comorbid distress of those referred for problematic gaming behaviours to better describe one's profile. Advancements in treatment for disordered gaming have been made, specifically with promising results in cognitive behavioral therapy (CBT) based psychotherapy (Li et al., 2020; Zajac et al., 2020). However, CBT-based psychotherapies targeting IGD often include exclusively cognitive techniques exclusively targeting depression (e.g., behavioural activation, or scheduling pleasurable non-gaming activities) or anxiety reduction techniques (e.g., systematic desensitization; Zajac et al., 2020). Therefore, programs and interventions designed for internet gaming disorder may benefit from including therapeutic techniques that target concurrently depression and anxiety.

Limitations and Future Research

It is important to note the present study presented a limited perspective regarding what encompasses distress. While it is true

Figure 2

Here We See Raw DAS Scores Discriminated by Distress Measure (i.e., Depression, Anxiety, Stress) and Latent Profile




Note. The horizontal dashed lines indicate cut-off scores for mild, moderate, severe, and extremely severe distress scores. Depression scores are classified as normal (0–4), mild (5–6), moderate (7–10), severe (11–13), and extremely severe (14+); anxiety scores are classified as normal (0–3), mild (4–5), moderate (6–7), severe (8–9), and extremely severe (10+); stress scores are classified as normal (0–7), mild (8–9), moderate (10–12), severe (13–16), and extremely severe (17+). Finally, we joined the three latent profiles with a regression line to highlight differences across profiles. DAS = Depression, Anxiety and Stress. See the online article for the color version of this figure.

that depression, anxiety, and stress have been postulated to possess a causal relationship with IGD (Griffiths, 2005; Griffiths et al., 2017; Kardefelt-Winther, 2014), the inclusion of only three broad indicators may fail to capture the scope of distress variability within the gaming population. Although not associated in a causal manner, incorporating symptoms positively associated with both distress and IGD (e.g., social withdrawal, fatigue, sleep disruption, Achab et al., 2011) may better describe the variety of distress experienced by the gamer and add more detail to potential risk profiles. Additionally, it

must be emphasized that there is a lack of clear temporal association between psychiatric features and the onset of gaming disorder (Balhara et al., 2018). Therefore, further research is required to better understand a potential network of factors influencing risk factors such as depression, anxiety, or stress and the abuse of online applications, such as gaming (Kardefelt-Winther, 2014; Zarate et al., 2022). Furthermore, the lack of longitudinal data poses an issue for the validity of these profiles over time. In this context, it is possible that the different distress profiles are constituting different stages of

Figure 3

DAS as a Function of IGD



Note. This plot shows relationships between IGD and distress symptoms (i.e., depression, anxiety, and stress) discriminated by distress profiles. As seen here, elevated levels of comorbid distress are associated with higher IGD scores. DAS = Depression, Anxiety and Stress; IGD = Internet Gaming Disorder. See the online article for the color version of this figure.

distress development for some individuals. Future research may wish to longitudinally profile gamers, using a broader range of distress symptoms or other factors thought to contribute to the development of IGD.

Conclusion

Findings, particularly those relating to IGD levels within distress profiles, have significant implications regarding the prevention, intervention, and assessment of IGD. Gamers presenting with higher levels of depression, anxiety, and stress should be considered as at greater risk of developing IGD. Thus, highlighting the need to target these individuals in primary care. Additionally, these findings call attention to the necessity of an efficacious treatment targeting depression, anxiety, and stress concurrently with IGD.

Résumé

Le trouble du jeu sur Internet (TJI) se caractérise par un engagement persistant et récurrent dans les jeux vidéo, à l'exclusion de toute autre activité, qui ne peut être contrôlé et qui entraîne des troubles importants du fonctionnement quotidien. Des recherches antérieures suggèrent que le trouble du jeu sur Internet est vécu différemment selon le profil du joueur, tandis que des symptômes de détresse tels que la dépression, l'anxiété et le stress ont été indépendamment associés au développement du trouble du jeu sur Internet. Il est intéressant de noter qu'aucune étude à ce jour n'a cherché à établir le profil des joueurs en fonction de ces trois psychopathologies. La présente étude visait à (a) établir le profil des joueurs concernant leur dépression, leur anxiété et leur stress, et (b) examiner les différences de niveaux de TJI entre les différents profils de détresse. Un échantillon composé de 968 joueurs (18 à 64 ans, Âge M = 29,54) a été évalué à l'aide de l'échelle de dépression, d'anxiété et de stress (DASS-21) et de l'échelle de troubles liés aux jeux sur Internet (IGDS9-SF). L'analyse des classes latentes (ACL) a permis d'identifier trois profils de détresse distincts. Ceux-ci englobent les joueurs à « comorbidité avec niveau de détresse élevé » (HDC; 25,9 %), « comorbidité avec niveau de détresse moyen » (MDC; 48,7 %) et « comorbidité avec niveau de détresse faible » (LDC; 25,4 %). Comme prévu, les profils de comorbidité de détresse plus élevés sont liés à des niveaux de TJI plus élevés. Les résultats suggèrent qu'il existe des profils de détresse différents selon la gravité des symptômes, les joueurs les plus en détresse présentant des niveaux de TJI plus élevés, indépendamment du symptôme de détresse spécifique qu'ils ont signalé. Ainsi, le diagnostic différentiel des symptômes de dépression, d'anxiété et de stress n'est peut-être pas nécessaire dans le contexte du traitement de TJI, mais les comportements de détresse devraient plutôt être ciblés simultanément et indépendamment de leur nature.

Mots-clés : anxiété, dépression, trouble du jeu sur Internet, analyse de classe latente, stress

References

Achab, S., Nicolier, M., Mauny, F., Monnin, J., Trojak, B., Vandel, P., Sechter, D., Gorwood, P., & Haffen, E. (2011). Massively multiplayer online role-playing games: Comparing characteristics of addict vs non-addict online recruited gamers in a French adult population.

- BMC Psychiatry*, 11(1), Article 144. <https://doi.org/10.1186/1471-244X-11-144>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). <https://doi.org/10.1176/appi.books.9780890425596>
- Anderson, E., Steen, E., & Stavropoulos, V. (2017). Internet use and problematic internet use: A systematic review of longitudinal research trends in adolescence and emergent adulthood. *International Journal of Adolescence and Youth*, 22(4), 430–454. <https://doi.org/10.1080/02673843.2016.1227716>
- Balhara, Y. P. S., Garg, H., Kumar, S., & Bhargava, R. (2018). Gaming disorder as a consequence of attempt at self-medication: Empirical support to the hypothesis. *Asian Journal of Psychiatry*, 31, 98–99. <https://doi.org/10.1016/j.ajp.2018.02.013>
- Billieux, J., Thorens, G., Khazaal, Y., Zullino, D., Achab, S., & Van der Linden, M. (2015). Problematic involvement in online games: A cluster analytic approach. *Computers in Human Behavior*, 43, 242–250. <https://doi.org/10.1016/j.chb.2014.10.055>
- Billieux, J., Schimmenti, A., Khazaal, Y., Maurage, P., & Heeren, A. (2015). Are we overpathologizing everyday life? A tenable blueprint for behavioral addiction research. *Journal of Behavioral Addictions*, 4(3), 119–123. <https://doi.org/10.1556/2006.4.2015.009>
- Brand, M., Young, K. S., Laier, C., Wölfling, K., & Potenza, M. N. (2016). Integrating psychological and neurobiological considerations regarding the development and maintenance of specific internet-use disorders: An Interaction of Person-Affect-Cognition-Execution (I-PACE) model. *Neuroscience and Biobehavioral Reviews*, 71, 252–266. <https://doi.org/10.1016/j.neubiorev.2016.08.033>
- Burleigh, T. L., Stavropoulos, V., Liew, L. W. L., Adams, B. L. M., & Griffiths, M. D. (2018). Depression, Internet Gaming Disorder, and the moderating effect of the gamer-avatar relationship: An exploratory longitudinal study. *International Journal of Mental Health and Addiction*, 16(1), 102–124. <https://doi.org/10.1007/s11469-017-9806-3>
- Celeux, G., & Soromenho, G. (1996). An entropy criterion for assessing the number of clusters in a mixture model. *Journal of Classification*, 13(2), 195–212. <https://doi.org/10.1007/BF01246098>
- Clark, S. L., & Muthén, B. (2009). *Relating latent class analysis results to variables not included in the analysis*. https://www.researchgate.net/publication/237346694_Relating_Latent_Class_Analysis_Results_to_Variables_not_Included_in_the_Analysis
- Coker, A. O., Coker, O. O., & Sanni, D. (2018). Psychometric properties of the 21-item Depression Anxiety Stress Scale (DASS-21). *African Research Review*, 12(2), Article 135. <https://doi.org/10.4314/afrrrev.v12i2.13>
- Colder Carras, M., & Kardefelt-Winther, D. (2018). When addiction symptoms and life problems diverge: A latent class analysis of problematic gaming in a representative multinational sample of European adolescents. *European Child & Adolescent Psychiatry*, 27(4), 513–525. <https://doi.org/10.1007/s00787-018-1108-1>
- Darvesh, N., Radhakrishnan, A., Lachance, C. C., Nincic, V., Sharpe, J. P., Ghassemi, M., Straus, S. E., & Tricco, A. C. (2020). Exploring the prevalence of gaming disorder and Internet gaming disorder: A rapid scoping review. *Systematic Reviews*, 9(1), Article 68. <https://doi.org/10.1186/s13643-020-01329-2>
- Davis, R. A. (2001). A cognitive-behavioral model of pathological Internet use. *Computers in Human Behavior*, 17(2), 187–195. [https://doi.org/10.1016/S0747-5632\(00\)00041-8](https://doi.org/10.1016/S0747-5632(00)00041-8)
- Faulkner, G., Irving, H., Adlaf, E. M., & Turner, N. (2015). Subtypes of adolescent video gamers: A latent class analysis. *International Journal of Mental Health and Addiction*, 13(1), 1–18. <https://doi.org/10.1007/s11469-014-9501-6>
- Goeders N. E. (2003). The impact of stress on addiction. *European neuropsychopharmacology: the journal of the European College of Neuropsychopharmacology*, 13(6), 435–441. [https://doi.org/10.1016/S0924-977X\(03\)90003-4](https://doi.org/10.1016/S0924-977X(03)90003-4)

- Goh, B. (2021, August 31). Three hours a week: Play time's over for China's young video gamers. *Reuters*. Retrieved October 6, 2021, from <https://www.reuters.com/world/china/china-rolls-out-new-rules-minors-online-gaming-xinhua-2021-08-30/>
- González-Bueso, V., Santamaría, J. J., Fernández, D., Merino, L., Montero, E., & Ribas, J. (2018). Association between Internet Gaming Disorder or pathological video-game use and comorbid psychopathology: A comprehensive review. *International Journal of Environmental Research and Public Health*, 15(4), Article E668. <https://doi.org/10.3390/ijerph15040668>
- Granic, I., Lobel, A., & Engels, R. C. M. E. (2014). The benefits of playing video games. *American Psychologist*, 69(1), 66–78. <https://doi.org/10.1037/a0034857>
- Griffiths, M. (2005). A “components” model of addiction within a biopsychosocial framework. *Journal of Substance Use*, 10(4), 191–197. <https://doi.org/10.1080/14659890500114359>
- Griffiths, M. D., Kuss, D. J., & Gortari, O. (2017). Videogames as therapy: An updated selective review of the medical and psychological literature. *International Journal of Privacy and Health Information Management*, 5(2), 71–96. <https://doi.org/10.4018/IJPHIM.2017070105>
- Ho, R. C., Zhang, M. W. B., Tsang, T. Y., Toh, A. H., Pan, F., Lu, Y., Cheng, C., Yip, P. S., Lam, L. T., Lai, C. M., & Mak, K. K. (2014). The association between internet addiction and psychiatric co-morbidity: A meta-analysis. *BMC Psychiatry*, 14(1), Article 183. <https://doi.org/10.1186/1471-244X-14-183>
- Kardefelt-Winther, D. (2014). A conceptual and methodological critique of internet addiction research: Towards a model of compensatory internet use. *Computers in Human Behavior*, 31, 351–354. <https://doi.org/10.1016/j.chb.2013.10.059>
- Kardefelt-Winther, D., Heeren, A., Schimmenti, A., van Rooij, A., Maurage, P., Carras, M., Edman, J., Blaszczynski, A., Khazaal, Y., & Billieux, J. (2017). How can we conceptualize behavioural addiction without pathologizing common behaviours? *Addiction*, 112(10), 1709–1715. <https://doi.org/10.1111/add.13763>
- Khantzian, E. J. (1997). The self-medication hypothesis of substance use disorders: A reconsideration and recent applications. *Harvard Review of Psychiatry*, 4(5), 231–244. <https://doi.org/10.3109/10673229709030550>
- Kim, B. N., & Ko, H. (2020). Psychometric properties of the Nine-Item Korean Internet Gaming Disorder Scale: Short form. *Cyberpsychology, Behavior and Social Networking*, 23(12), 854–859. <https://doi.org/10.1089/cyber.2020.0227>
- Larose, C., Harel, O., Kordas, K., & Dey, D. K. (2016). Latent class analysis of incomplete data via an entropy-based criterion. *Statistical Methodology*, 32, 107–121. <https://doi.org/10.1016/j.stamet.2016.04.004>
- Le, M. T. H., Tran, T. D., Holton, S., Nguyen, H. T., Wolfe, R., & Fisher, J. (2017). Reliability, convergent validity and factor structure of the DASS-21 in a sample of Vietnamese adolescents. *PLOS ONE*, 12(7), Article e0180557. <https://doi.org/10.1371/journal.pone.0180557>
- Lee, S.-Y., Lee, H. K., & Choo, H. (2017). Typology of Internet gaming disorder and its clinical implications. *Psychiatry and Clinical Neurosciences*, 71(7), 479–491. <https://doi.org/10.1111/pcn.12457>
- Lemmens, J. S., Valkenburg, P. M., & Gentile, D. A. (2015). The Internet Gaming Disorder Scale. *Psychological Assessment*, 27(2), 567–582. <https://doi.org/10.1037/pas0000062>
- Li, J., Li, X., Jiang, J., Xu, X., Wu, J., Xu, Y., Lin, X., Hall, J., Xu, H., Xu, J., & Xu, X. (2020). The effect of cognitive behavioral therapy on depression, anxiety, and stress in patients with COVID-19: A randomized controlled trial. *Frontiers in Psychiatry*, 11, Article 580827. <https://doi.org/10.3389/fpsy.2020.580827>
- Liew, L. W. L., Stavropoulos, V., Adams, B. L. M., Burleigh, T. L., & Griffiths, M. D. (2018). Internet Gaming Disorder: The interplay between physical activity and user–avatar relationship. *Behaviour & Information Technology*, 37(6), 558–574. <https://doi.org/10.1080/0144929X.2018.1464599>
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck depression and anxiety inventories. *Behaviour Research and Therapy*, 33(3), 335–343. [https://doi.org/10.1016/0005-7967\(94\)00075-U](https://doi.org/10.1016/0005-7967(94)00075-U)
- Männikkö, N., Ruotsalainen, H., Miettunen, J., Pontes, H. M., & Kääriäinen, M. (2020). Problematic gaming behaviour and health-related outcomes: A systematic review and meta-analysis. *Journal of Health Psychology*, 25(1), 67–81. <https://doi.org/10.1177/1359105317740414>
- Martín-Fernández, M., Matalí, J. L., García-Sánchez, S., Pardo, M., Lleras, M., & Castellano-Tejedor, C. (2016). Adolescents with Internet Gaming Disorder (IGD): Profiles and treatment response. *Addictions*, 29(2), 125–133. <https://doi.org/10.20882/adicciones.890>
- Masyn, K. E. (2013). Latent class analysis and finite mixture modeling: Statistical analysis. In T. D. Little (Ed.), *The Oxford handbook of quantitative methods* (pp. 551–611). Oxford University Press.
- McLachlan, G. J. (1987). On bootstrapping the likelihood ratio test statistic for the number of components in a normal mixture. *Applied Statistics*, 36(3), 318–324. <https://doi.org/10.2307/2347790>
- Muthén, L. K., & Muthén, B. (2016). *Mplus. The comprehensive modelling program for applied researchers. User's guide*, 5.
- O'Brien, C. (2018). *A systematic review and meta-analysis of player motivations and problematic involvement in multiplayer online games: Exploring an alternative diagnostic approach that minimizes the risk of pathologizing healthy gaming behaviors* (Order No. 10974517). Available from ProQuest One Academic. (2129737608). <https://www.proquest.com/dissertations-theses/systematic-review-meta-analysis-player/docview/2129737608/se-2?accountid=14844>
- Ostinelli, E. G., Zangani, C., Giordano, B., Maestri, D., Gambini, O., D'Agostino, A., Furukawa, T. A., & Purgato, M. (2021). Depressive symptoms and depression in individuals with internet gaming disorder: A systematic review and meta-analysis. *Journal of Affective Disorders*, 284, 136–142. <https://doi.org/10.1016/j.jad.2021.02.014>
- Pontes, H. M., Király, O., Demetrovics, Z., & Griffiths, M. D. (2014). The conceptualisation and measurement of DSM-5 Internet Gaming Disorder: The development of the IGD-20 Test. *PLOS ONE*, 9(10), Article e110137. <https://doi.org/10.1371/journal.pone.0110137>
- Raith, L., Bignill, J., Stavropoulos, V., Millea, P., Allen, A., Stallman, H. M., Mason, J., De Regt, T., Wood, A., & Kannis-Dymand, L. (2021). Massively multiplayer online games and well-being: A systematic literature review. *Frontiers in Psychology*, 12, Article 698799. <https://doi.org/10.3389/fpsyg.2021.698799>
- Rosenberg, J., Beymer, P., Anderson, D., van Lissa, C., & Schmidt, J. (2019). tidyLPA: An R package to easily carry out Latent Profile Analysis (LPA) using open-source or commercial software. *Journal of Open Source Software*, 3(30), Article 978. <https://doi.org/10.21105/joss.00978>
- Schafer, J. L. (1999). Multiple imputation: A primer. *Statistical Methods in Medical Research*, 8(1), 3–15. <https://doi.org/10.1177/096228029900800102>
- Stavropoulos, V., Beard, C., Griffiths, M. D., Buleigh, T., Gomez, R., & Pontes, H. M. (2018). Measurement invariance of the Internet Gaming Disorder Scale–Short-Form (IGDS9-SF) between Australia, the USA, and the UK. *International Journal of Mental Health and Addiction*, 16(2), 377–392. <https://doi.org/10.1007/s11469-017-9786-3>
- Stavropoulos, V., Anderson, E. E., Beard, C., Latifi, M. Q., Kuss, D., & Griffiths, M. (2018). A preliminary cross-cultural study of Hikikomori and Internet Gaming Disorder: The moderating effects of game-playing time and living with parents. *Addictive Behaviors Reports*, 9, Article 100137. <https://doi.org/10.1016/j.abrep.2018.10.001>
- Stavropoulos, V., Gomez, R., Mueller, A., Yucel, M., & Griffiths, M. (2020). User-avatar bond profiles: How do they associate with disordered gaming? *Addictive Behaviors*, 103, Article 106245. <https://doi.org/10.1016/j.addbeh.2019.106245>

- Stavropoulos, V., Motti-Stefanidi, F., & Griffiths, M. D. (2021). Risks and opportunities for youth in the digital era: A cyber-developmental approach to mental health. *European Psychologist*. Advance online publication. <https://doi.org/10.1027/1016-9040/a000451>
- Stavropoulos, V., Vassallo, J., Burleigh, T. L., Gomez, R., & Carras, M. C. (2021). The role of internet gaming in the association between anxiety and depression: A preliminary cross-sectional study. *Asia-Pacific Psychiatry: Official Journal of the Pacific Rim College of Psychiatrists*, Article e12474. <https://doi.org/10.1111/appy.12474>
- Stavropoulos, V., Gomez, R., & Griffiths, M. D. (2021). In search of the optimum structural model for Internet Gaming Disorder. *BMC Psychiatry*, 21(1), Article 176. <https://doi.org/10.1186/s12888-021-03148-8>
- Tullett-Prado, D., Stavropoulos, V., Mueller, K., Sharples, J., & Footitt, T. A. (2021). Internet Gaming Disorder profiles and their associations with social engagement behaviours. *Journal of Psychiatric Research*, 138, 393–403. <https://doi.org/10.1016/j.jpsychires.2021.04.037>
- Ünöl, H., Koç, A. Ş., Sayar, G. H., Stavropoulos, V., Kircaburun, K., & Griffiths, M. D. (2020). Measurement, profiles, prevalence, and psychological risk factors of problematic gaming among the Turkish community: A large-scale national study. *International Journal of Mental Health and Addiction*, 19(5), 1662–1682. <https://doi.org/10.1007/s11469-020-00254-8>
- Valentino, R. J., Lucki, I., & Van Bockstaele, E. (2010). Corticotropin-releasing factor in the dorsal raphe nucleus: Linking stress coping and addiction. *Brain Research*, 1314, 29–37. <https://doi.org/10.1016/j.brainres.2009.09.100>
- van Rooij, A. J., Ferguson, C. J., Colder Carras, M., Kardefelt-Winther, D., Shi, J., Aarseth, E., Bean, A. M., Bergmark, K. H., Brus, A., Coulson, M., Deleuze, J., Dullur, P., Dunkels, E., Edman, J., Elson, M., Etchells, P. J., Fiskaali, A., Granic, I., . . . Przybylski, A. K. (2018). A weak scientific basis for gaming disorder: Let us err on the side of caution. *Journal of Behavioral Addictions*, 7(1), 1–9. <https://doi.org/10.1556/2006.7.2018.19>
- Wong, H. Y., Mo, H. Y., Potenza, M. N., Chan, M. N. M., Lau, W. M., Chui, T. K., Pakpour, A. H., & Lin, C.-Y. (2020). Relationships between severity of Internet Gaming Disorder, severity of problematic social media use, sleep quality and psychological distress. *International Journal of Environmental Research and Public Health*, 17(6), Article E1879. <https://doi.org/10.3390/ijerph17061879>
- World Health Organization (2019). *International classification of diseases* (11th ed.).
- Yen, J.-Y., Lin, H.-C., Chou, W.-P., Liu, T.-L., & Ko, C.-H. (2019). Associations among Resilience, Stress, Depression, and Internet Gaming Disorder in young adults. *International Journal of Environmental Research and Public Health*, 16(17), Article E3181. <https://doi.org/10.3390/ijerph16173181>
- Zajac, K., Ginley, M. K., & Chang, R. (2020). Treatments of internet gaming disorder: A systematic review of the evidence. *Expert Review of Neurotherapeutics*, 20(1), 85–93. <https://doi.org/10.1080/14737175.2020.1671824>
- Zarate, D., Ball, M., Montag, C., Prokofieva, M., & Stavropoulos, V. (2022). Unravelling the web of addictions: A network analysis approach. *Addictive Behaviors Reports*, Article 100406. <https://doi.org/10.1016/j.abrep.2022.100406>

Received December 8, 2021

Revision received March 8, 2022

Accepted March 10, 2022 ■