

Can Reptiles Perceive Visual Illusions? Delboeuf Illusion in Red-Footed Tortoise (*Chelonoidis carbonaria*) and Bearded Dragon (*Pogona vitticeps*)

Maria Santacà
University of Padova

Maria Elena Miletto Petrazzini
Queen Mary University of London

Christian Agrillo
University of Padova

Anna Wilkinson
University of Lincoln

Optical illusions have been widely used to compare visual perception among vertebrates because they can reveal how the system is able to adapt to visual input. Sensitivity to visual illusions has never been studied in reptiles. Here, we investigated whether red-footed tortoises, *Chelonoidis carbonaria*, and bearded dragons, *Pogona vitticeps*, perceive the Delboeuf illusion. This illusion involves the misperception of the size of a target circle depending upon the context in which it is presented. We adopted the same size discrimination for both species to compare their performance. Animals were presented with two different types of trials. In control trials, they received two different-sized food portions on two plates of the same size. In test trials, they received two same-sized food portions but presented on two different-sized plates. If they perceived the illusion in the same way as humans, we expected them to select the food portion presented on the smaller plate. The tortoises exhibited poor performance in the control trials, which prevented us from drawing any conclusions about their perception of the Delboeuf illusion. In contrast, the bearded dragons selected the larger amount of food in control trials. In test trials, they selected the portion presented on the smaller plate significantly more often than chance, suggesting a human-like sensitivity to the Delboeuf illusion. Our study provides the first evidence of the perception of a visual illusion in a reptile species, suggesting that rather than simply detecting visual input, they interpret sensory information captured by photoreceptors.

Keywords: reptile cognition, visual illusions, size illusions, Delboeuf illusion

Animals are found in almost every habitat on earth, and they have developed extraordinarily different eyes and ways of seeing

the world (Lazareva, Shimizu, & Wasserman, 2012). As such, it is essential to investigate similarities and differences in perception between species. Optical illusions, subjective interpretations that differ from physical stimulation, are becoming an increasingly popular tool to investigate animal visual perception (Feng, Chouinard, Howell, & Bennett, 2017; Kelley & Kelley, 2014). This approach allows us to assess whether animals interpret visual inputs as humans do or whether they detect visual inputs with little or no variability (Feng et al., 2017). Investigation of species differences in susceptibility to illusions may also shed light on the impact of environmental and evolutionary pressures on visual perception (Feng et al., 2017).

The majority of the current work in this area has focused on mammals, such as chimpanzees (*Pan troglodytes*; Fujita, 1997; Parrish & Beran, 2014), rhesus monkeys (*Macaca mulatta*; Agrillo, Parrish, & Beran, 2014a, 2014b; Bayne & Davis, 1983; Fujita, 1997), capuchin monkeys (*Cebus paella*; Agrillo et al., 2014b; Parrish, Agrillo, Perdue, & Beran, 2016; Suganuma, Pessoa, Monge-Fuentes, Castro, & Tavares, 2007), baboons (*Papio papio*; Parron & Fagot, 2007), ring-tailed lemurs (*Lemur catta*; Santacà, Regaiolli, Miletto Petrazzini, Spiezio, & Agrillo, 2017), dogs [*Canis familiaris*; Byosiére et al., 2017; Keep, Zulch, & Wilkinson, 2018; Miletto Petrazzini, Bisazza, & Agrillo, 2017], and bottlenose dolphins (*Tursiops truncatus*; Murayama, Usui, Takeda, Kato, & Maejima, 2012). Only three

This article was published Online First March 21, 2019.

Maria Santacà, Department of General Psychology, University of Padova; Maria Elena Miletto Petrazzini, School of Biological and Chemical Science, Queen Mary University of London; Christian Agrillo, Department of General Psychology, University of Padova; Anna Wilkinson, School of Life Sciences, University of Lincoln.

We thank the University of Lincoln Minster House technical team, in particular Paul Sheridan, for their help with animal care and setup. This study was supported by “STARS@unipd” (ANIM_ILLUS) grant from the University of Padova to Christian Agrillo and by Marie Skłodowska-Curie individual fellowship (SBCN1E4R) from the European Union’s Horizon 2020 to Maria Elena Miletto Petrazzini. We also thank all the students that helped with the video analysis. This research was approved by the ethics committee of the School of Life Sciences, University of Lincoln (CoSREC364). Applicable national guidelines for the care and use of animals were followed. The present work has been done in the scope of the research grant ‘Dipartimenti di Eccellenza’ use inspired basic research entitled: ‘Innovative methods or technologies for assessment, intervention, or enhancement of psychological functions (cognitive, emotional or behavioural)’.

Correspondence concerning this article should be addressed to Maria Santacà, Department of General Psychology, University of Padova, Via Venezia 8, 35131 Padova, Italy. E-mail: santacamaria@gmail.com

species of birds have been investigated regarding their susceptibility to illusory patterns: gray parrot (*Psittacus erithacus*; Pepperberg & Nakayama, 2016; Pepperberg, Vicinay, & Cavanagh, 2008), domestic chicks (*Gallus gallus*; Rosa Salva, Rugani, Cavazzana, Regolin, & Vallortigara, 2013; Watanabe, Nakamura, & Fujita, 2013), and pigeons (*Columba livia*; Nakamura, Watanabe, & Fujita, 2008; Watanabe, Nakamura, & Fujita, 2011). In elasmobranchs, only bamboo sharks have been studied (*Chiloscyllium griseum*; Fuss, Bleckmann, & Schlüssel, 2014; Fuss & Schlüssel, 2017), whereas a larger range of teleost fishes has been studied, including zebrafish (*Danio rerio*; Gori, Agrillo, Dadda, & Bisazza, 2015), guppies (*Poecilia reticulata*; Agrillo, Miletto Petrazzini, & Bisazza, 2016; Gori et al., 2015), redbellied splitfin fish (*Xenotoca eiseni*; Sovrano, Albertazzi, & Rosa Salva, 2015; Sovrano & Bisazza, 2009; Sovrano, Da Pos, & Albertazzi, 2016), goldfish (*Carassius auratus*; Wyzisk & Neumeyer, 2007), and damselfish (*Chromis chromis*; Fuss & Schlüssel, 2017). Susceptibility to illusory patterns was also studied in invertebrates, such as the perception of illusory contours and contextual size illusions by honeybees (*Apis mellifera*; Horridge, Zhang, & O'Carroll, 1992; Howard, Avarguès-Weber, Garcia, Stuart-Fox, & Dyer, 2017). Some of these species showed a human-like perception of illusory phenomena, whereas others exhibited a reversed illusion, meaning that animals perceived a sort of illusion but in the opposite way to human observers (for example, pigeons [Nakamura et al., 2008; Watanabe et al., 2011] and bantams [Watanabe et al., 2013]). Furthermore, some species did not appear to perceive the illusion (for example, rhesus monkeys [Agrillo et al., 2014b] and bamboo sharks [Fuss et al., 2014]). However, in some cases there is evidence that different methods of investigating illusion sensitivity can lead to different results in the same species. One remarkable example is the case of bantam chickens and the Ebbinghaus illusion, a relative size perception illusion. Rosa Salva et al. (2013) demonstrated that 4-day-old chicks perceived the illusion in a human-like direction, whereas Nakamura, Watanabe, and Fujita (2014) concluded the opposite testing 6-month-old chickens.

At this stage, no firm conclusions can be made regarding the universality of perceptual mechanisms, as many species are yet to be studied. One specific class, the Reptilia, has not been investigated. Reptiles were long considered to be sluggish and unintelligent; however, when tested under appropriate experimental conditions, they exhibit an impressive array of cognitive abilities (for example, red-footed tortoise, *Chelonoidis carbonaria*, and bearded dragon, *Pogona vitticeps*; Kis, Huber, & Wilkinson, 2015; Mueller-Paul et al., 2014; reviewed by Matsubara, Deeming, & Wilkinson, 2017; Wilkinson & Glass, 2018). Work has recently shown that these two species are able to perceive similarities between pictures and the objects that they represent (Wilkinson, Mueller-Paul, & Huber, 2013), respond to video stimuli (Siviter, Deeming, van Giezen, & Wilkinson, 2018; Wilkinson, Sebanz, Mandl, & Huber, 2011), and, crucially for this experiment, discriminate between different quantities of food reward (and remember this association for 18 months; Soldati, Burman, John, Pike, & Wilkinson, 2017). This ability was also found in another reptile species, the Italian wall lizard, *Podarcis sicula* (Miletto Petrazzini, Fraccaroli, et al., 2017).

This study therefore investigated how two reptile species, the red-footed tortoise (*Chelonoidis carbonaria*) and the bearded dragon (*Pogona vitticeps*), perceive one of the most famous geometrical illusions, the Delboeuf illusion. In the most familiar version of this illusion, two identical circles are near each other.




Figure 1. Delboeuf illusion. This illusion occurs when two same-sized circles are perceived to be different depending on the context in which they are presented. In the classical version, the central target circles are physically identical, but human observers typically underestimate the size of the one included in the larger ring and tend to overestimate the size of the one encompassed by the smaller ring.

One is encircled by a circle with a small circumference, whereas the other is encircled by a circle with a large circumference (Figure 1). Humans tend to perceive the former as larger than the latter despite them being the same size (Pressey, 1977). Similarly, capuchin monkeys (Parrish, Brosnan, & Beran, 2015) have been shown to overestimate the size of a circle when surrounded by a circle with a small circumference. In another version of the illusion, used with chimpanzees, the target circles were replaced by food portions. Chimpanzees (Parrish & Beran, 2014) tended to overestimate the dimension of a food portion included in the smaller array, whereas they tended to underestimate the dimension of the same food portion when it was presented in the larger array. Interestingly it has been demonstrated that humans' perception of food portion is influenced by the Delboeuf pattern. Humans overestimate food sizes when food is presented on small plates (Davis, Payne, & Bui, 2016; Van Ittersum & Wansink, 2007; Wansink & Cheney, 2005). In humans, the illusion seems to be a combination of both assimilation and contrast effects (King, 1988). The food portion presented on the smaller plate is thought to assimilate to the contour of the plate, leading to it being perceived as larger than it is, whereas the food portion presented on the larger plate is thought to contrast to the contour of the plate, leading to an underestimation of the food portion size. To reduce the methodological variability and make a reliable comparison between the different species, we adapted the same spontaneous choice procedure adopted in the studies of chimpanzees (Parrish & Beran, 2014), lemurs (Santacà et al., 2017), and dogs (Miletto Petrazzini, Bisazza, & Agrillo, 2017) to investigate this question in reptiles. In spontaneous choice tests, animals are thought to exhibit their natural behavior, and their performance is likely to reflect the cognitive and perceptual functions they would activate in nature. On the contrary, intensive training procedures might lead to extraordinary performances through experience and the recruitment of other neural networks to accommodate for the extensive requirements of a specific cognitive task (Agrillo & Bisazza, 2014).

Reptiles were observed spontaneously selecting one of two arrays containing food portions. We arranged two different control trials, to verify the tendency of the subjects to maximize the food

intake in our experimental context, and test trials with the illusory pattern. In the control trials, we presented two different-sized food portions (with a ratio of 0.67 between them, a ratio commonly used in spontaneous discrimination tasks with animals, including reptiles; Bánszegi, Urrutia, Szenczi, & Hudson, 2016; Miletto Petrazzini, Fraccaroli, et al., 2017; Miletto Petrazzini & Wynne, 2016) on two same-sized arrays. In test trials, we presented two identical food portions but on two different-sized arrays, one small and one large. If red-footed tortoises and bearded-dragons perceived the Delboeuf illusion in a human-like way, they were expected to choose the portion presented on the smaller array.

Materials and Method

Subjects

A total of eight red-footed tortoises (*Chelonoidis carbonaria*; five females and three males) and 12 bearded-dragons (*Pogona vitticeps*; eight females and four males), participated in this study (Tables 1 and 2). The reptiles were maintained at the School of Life Sciences, University of Lincoln. The tortoises were housed all together in a heated (28°C) and humidified room, whereas the bearded dragons were housed individually or in pairs in vivaria in an adjacent room. All animals had permanent access to fresh water, shelter, ultraviolet light, and heat lamps. No subject was food deprived during the experiment, but they received a favored food reward during the experiment. All animals were handled by humans on a daily basis. None of the subjects were experimentally naïve (Moszuti, Wilkinson, & Burman, 2017; Siviter et al., 2017, 2018; Soldati et al., 2017), but they had not previously taken part in tests to investigate the susceptibility to visual illusions. Each subject was tested individually.

Stimuli and Apparatus

The stimuli consisted of mango jelly for the tortoises and vegetable extract (kale, cucumber, and mint) jelly for the bearded dragons, highly preferred food for both species. The jellies were prepared each day and cut with a circular cutter to get a consistent round form. Then, each jelly was placed in the middle of a 7.5 × 8.5 cm black plastic card. In each card, one central white circle

represented the plate. Two different sizes of plates were used: The larger plates had a diameter of 4.92 cm, whereas the smaller plates had a diameter of 1.83 cm. Two different portion sizes were presented to the reptiles: The larger food portion was 1.5 cm in diameter (area = 1.77 cm²), whereas the smaller food portion was 1.23 cm in diameter (area = 1.19 cm²). Each card was presented on an L-shaped steel bracket (7.5 × 8.5 × 4 cm) to improve the visibility of the arrays from subjects' points of view.

The experiments were run in an arena measuring 100 × 100 cm (Figure 2) located in a room maintained at 28°C (±3°C). To ensure the animals could see the entire array, they were positioned at the top of a ramp inclined by a 36.02° angle with both food plates at the bottom. The same apparatus was used for both species with two differences. For the tortoises, the inner part of the arena was covered with grip and dark bark only in the choice area, whereas for the bearded dragons it was entirely covered with black plastic. The arena was also covered with a wire mesh for the bearded dragons to avoid the possibility of their escaping from the apparatus. To reduce the possibility of subjects using olfactory cues, the apparatus was cleaned after each trial.

Procedure

Before starting the test phase, each animal received a short familiarization phase. During the familiarization phase, subjects were presented with a single card containing some pieces of jelly in the center of the choice area. This procedure allowed us to habituate the animals to the cards and to ensure they ate the jelly in the choice area. All animals did this readily, and therefore this phase lasted for only one day; each animal receive a total of eight trials. The test phase began as soon as the familiarization phase was complete. For the illusory investigation, reptiles received 12 sessions, each consisting of four trials, receiving a total of 48 trials. They received two sessions each day with at least a 1-hr interval between sessions. Both control and test trials were presented to the subjects (Figure 3); control trials were used to assess their motivation to choose the larger food portion. In half of the control trials (Control A), the two different-sized portions of jelly were both on small plates, whereas in the other half (Control B), they were presented on large plates. In these trials, the physical difference between the two portions was equal to 0.67 (Bánszegi et al., 2016;

Table 1
Information of the Bearded Dragons Participating in the Study and Individual Performance

Subject	Control A	Control B	Control C	Test trials
Shuriken	12/16, $\chi^2 = 4.000$, $p = .047^*$	12/16, $\chi^2 = 4.000$, $p = .047^*$	12/16, $\chi^2 = 4.000$, $p = .047^*$	12/16, $\chi^2 = 4.000$, $p = 0.047^*$
Malie	11/16, $\chi^2 = 2.250$, $p = .134$	12/16, $\chi^2 = 4.000$, $p = .047^*$	11/16, $\chi^2 = 2.250$, $p = .134$	12/16, $\chi^2 = 4.000$, $p = 0.047^*$
Nimoy	6/16, $\chi^2 = 1.000$, $p = .317$	9/16, $\chi^2 = 0.250$, $p = .617$	11/16, $\chi^2 = 2.250$, $p = .134$	11/16, $\chi^2 = 2.250$, $p = 0.134$
Quadra	12/16, $\chi^2 = 4.000$, $p = .047^*$	10/16, $\chi^2 = 1.000$, $p = .317$	14/16, $\chi^2 = 9.000$, $p = .003^*$	8/16, $\chi^2 = .000$, $p = 1.000$
Norbert	12/16, $\chi^2 = 4.000$, $p = .047^*$	10/16, $\chi^2 = 1.000$, $p = .317$	13/16, $\chi^2 = 6.250$, $p = .012^*$	13/16, $\chi^2 = 6.250$, $p = 0.012^*$
Dr. Tom Pike	11/16, $\chi^2 = 2.250$, $p = .134$	12/16, $\chi^2 = 4.000$, $p = .047^*$	13/16, $\chi^2 = 6.250$, $p = .012^*$	12/16, $\chi^2 = 4.000$, $p = 0.047^*$
Alberta	13/16, $\chi^2 = 6.250$, $p = .012^*$	11/16, $\chi^2 = 2.250$, $p = .134$	12/16, $\chi^2 = 4.000$, $p = .047^*$	11/16, $\chi^2 = 2.250$, $p = 0.134$
Heinz	14/16, $\chi^2 = 9.000$, $p = .003^*$	11/16, $\chi^2 = 2.250$, $p = .134$	N/A	13/16, $\chi^2 = 6.250$, $p = 0.012^*$
Cecilia	12/16, $\chi^2 = 4.000$, $p = .047^*$	11/16, $\chi^2 = 2.250$, $p = .134$	N/A	10/16, $\chi^2 = 1.000$, $p = 0.317$
Oscar	15/16, $\chi^2 = 12.250$, $p = .0004^*$	13/16, $\chi^2 = 6.250$, $p = .012^*$	13/16, $\chi^2 = 6.250$, $p = .012^*$	12/16, $\chi^2 = 4.000$, $p = 0.047^*$
Mushu	13/16, $\chi^2 = 6.250$, $p = .012^*$	11/16, $\chi^2 = 2.250$, $p = .134$	10/16, $\chi^2 = 1.000$, $p = .317$	9/16, $\chi^2 = 0.250$, $p = 0.617$
Haku	13/16, $\chi^2 = 6.250$, $p = .012^*$	13/16, $\chi^2 = 6.250$, $p = .012^*$	N/A	12/16, $\chi^2 = 4.000$, $p = 0.047^*$

Note. Control trials = frequency of choices for the larger quantity; test trials = frequency of choices for the portion of food on the smaller plate.

* denotes a significant departure from chance at χ^2 test ($p < .05$).

Table 2
Information of the Red-Footed Tortoises Participating in the Study and Individual Performance

Subject	Control A	Control B	Test trials
6 (Savina)	12/16, $\chi^2 = 4.000$, $p = 0.047^*$	8/16, $\chi^2 = 0.000$, $p = 1.000$	8/16, $\chi^2 = 0.000$, $p = 1.000$
7 (Patty)	8/16, $\chi^2 = 0.000$, $p = 1.000$	9/16, $\chi^2 = 0.250$, $p = 0.617$	7/16, $\chi^2 = 0.250$, $p = 0.617$
8 (Charles)	12/16, $\chi^2 = 4.000$, $p = 0.047^*$	6/16, $\chi^2 = 1.000$, $p = 0.317$	11/16, $\chi^2 = 2.250$, $p = 0.134$
9 (Mozart)	9/16, $\chi^2 = 0.250$, $p = 0.617$	8/16, $\chi^2 = 0.000$, $p = 1.000$	8/16, $\chi^2 = 0.000$, $p = 1.000$
10 (Seisue)	8/16, $\chi^2 = 0.000$, $p = 1.000$	8/16, $\chi^2 = 0.000$, $p = 1.000$	7/16, $\chi^2 = 0.250$, $p = 0.617$
19 (T19)	9/16, $\chi^2 = 0.250$, $p = 0.617$	10/16, $\chi^2 = 1.000$, $p = 0.317$	8/16, $\chi^2 = 0.000$, $p = 1.000$
24 (Ranieri)	3/16, $\chi^2 = 6.250$, $p = 0.012^*$	8/16, $\chi^2 = 0.000$, $p = 1.000$	8/16, $\chi^2 = 0.000$, $p = 1.000$
300 (Gerard)	9/16, $\chi^2 = 0.250$, $p = 0.617$	12/16, $\chi^2 = 4.000$, $p = 0.047^*$	9/16, $\chi^2 = 0.250$, $p = 0.617$

Note. Control trials = frequency of choices for the larger quantity; test trials = frequency of choices for the portion of food on the smaller plate.

* denotes a significant departure from chance at χ^2 test ($p < .05$).

Lucon-Xiccato, Miletto Petrazzini, Agrillo, & Bisazza, 2015; Miletto Petrazzini, Fraccaroli, et al., 2017; Miletto Petrazzini & Wynne, 2016). In contrast, in the test trials, the subjects were presented with two same-sized jelly portions, one on a large plate and the other on a small plate. The ratio between the area of the jelly and the smaller plate was equal to 0.67, a ratio commonly used in human research to elicit the Delboeuf illusion (Piaget, 1957). In total, each subject was presented with 16 trials for each condition. The sequence of presentation of the trials was pseudo-randomized with the restriction that the test trial was never presented more than twice in a row and a session never began with a test trial. The position (left/right) of the larger food portion in the control trials and of the plate size (large/small) in the test trials was counterbalanced across trials. Each session was recorded, and the videos were analyzed to note the reptiles' choices (defined as the first jelly touched by the animal and in the meantime the other jelly was removed). One third of the videos of both species were

analyzed by an additional observer, and interrater reliability was excellent (Pearson's correlation $r = 1.0$, $p < .0001$).

The results of this study left open the possibility that the bearded dragons learned to avoid as much white as possible. Indeed, in control trials, the larger portion of food was intrinsically encircled by a smaller portion of white plate; in test trials, the choice for the smaller plate was also characterized by the fact that a smaller portion of the plate was visible in the chosen-array compared with the other option. To assess whether the subjects had learnt the rule to choose the configuration with less white, we arranged an additional type of control trials (Control C), presenting the two different-sized food portions in identical backgrounds without plates (Figure 3c). We tested nine out of 12 bearded dragons (two had died of natural causes and one was unwell at the time of testing) adopting the same procedure of the previous tests with the exception of receiving only one type of trials: They received two sessions each day,




Figure 2. Experimental setup. Three-dimensional representation of the experimental apparatus. Subjects had to descend a ramp in order to reach the plates containing the jellies. As the dependent variable, we recorded the first food portion touched by the animal in each trial.




Figure 3. Stimuli. Different or equal-sized food portions were presented on white plates: (a) Control A (different food portions in two identical small plates); (b) Control B (different food portions in two identical large plates); (c) Control C (different food portions in identical neutral backgrounds); (d) test trials (equal-sized food portions in two different-sized plates).

each consisting of four trials, receiving a total of 16 trials of Control C. Therefore, the nine bearded dragons received a total of 64 trials divided in 16 trials for each condition. The sequence of presentation of the trials was randomized with the restriction that the larger food portion was never presented more than twice in a row in the same position (left/right).

Data Analysis

Statistical analyses were performed using IBM SPSS 24.0 (SPSS Inc., Chicago, IL). Individual analyses were performed on the frequency of choices of the larger food portion in the control trials and of the food portion presented on the smaller plate (the apparently larger portion from a human perspective) in the test trials. Two-tailed χ^2 tests ($\alpha = .05$) were performed. Data were also analyzed at population level. Not all data were normally distributed, and hence we used parametric statistics for the data normally distributed (Shapiro-Wilk test, $p > .05$) and nonparametric statistics for the data that were not normally distributed (Shapiro-Wilk test, $p < .05$). One-sample t tests (chance level = 0.5) were performed to assess the discrimination of the two food portions in control trials and whether reptiles selected the apparently larger food portion (to humans) more than chance in test trials. The performances in the two types of control trials were compared using paired t tests. To investigate a possible change in performance in all types of trials over sessions, we performed an analysis of variance.

Results

Pogona vitticeps

Individual analyses (χ^2 tests) on the frequency of choices for the larger food portion showed that nine bearded dragons out of 12 in Control A and five subjects in Control B significantly selected the larger food portion (Table 1). Group analysis showed a significant discrimination of the larger quantity in both Control A ($M = 0.750$, 95% confidence interval [CI] [0.662, 0.838]; one-sample Wilcoxon signed-ranks test $p = .003$) and Control B ($M = 0.703$, 95% CI [0.655, 0.703], $t(11) = 9.275$, $p < .001$). No difference was found between the two control trials (related-sample Wilcoxon signed-ranks test $p = .162$).

In the test trials, individual analyses showed that seven out of the 12 bearded dragons significantly selected the food portion presented on the smaller plate (the one apparently larger to humans; Table 1). Group analyses also showed a significant preference for the food portion presented on the smaller plate ($M = 0.703$, 95% CI [0.640, 0.765], $t(11) = 7.294$, $p < .001$; Figure 4a). The analysis of variance revealed a nonsignificant difference in the bearded dragons' performance as a function of time in all types of trials, $F_{(11, 12)} = 2.478$, $p = .0671$, indicating a lack of learning effect during the experiment.

In Control C trials, individual analyses (χ^2 tests) on the frequency of choices for the larger food portion showed that six out of nine bearded dragons significantly selected the larger food portion (Table 1). Group analysis showed a significant discrimination of the larger quantity ($M = 0.757$, 95% CI [0.696, 0.818], $t(8) = 9.7167$, $p < .01$; Figure 4a). To further exclude the possibility that they simply learned to avoid as much white as possible, we analyzed the performance in the first trial of Control C: Eight out of nine bearded dragons selected the bigger food portion the first time they were presented with such trial. This result thus shows that bearded dragons were not driven by a tendency to avoid as much white as possible, but instead they focused on the biologically relevant stimuli presented in the arrays, the food portions.




Figure 4. Results. The y axis refers to the proportion of choices for the larger food portion in the control trials and the proportion of choices for the food portion presented on the small plate in the test trials. Bearded dragons (a), but not tortoises (b), selected the larger food portion in all control trials. Tortoises did not show a preference for any portion in test trials (b), whereas bearded dragons significantly chose more than chance the food portion included in the small plate suggesting a human-like perception of the Delboeuf illusion (a). Bars represent the standard error. The asterisk (*) denotes a significant departure from chance level.

Chelonoidis carbonaria

Chi-square tests showed that only two tortoises significantly selected the larger food portion in Control A and only one in Control B (Table 2). Contrary to expectations, one subject significantly selected the smaller portion in Control A. Group analyses revealed the lack of significant preference for any portion size in either type of control trial: Control A, $M = 0.547$, 95% CI [0.340, 0.694]; one-sample t test $t(7) = 0.753$, $p = .476$, and Control B, $M = 0.539$, 95% CI [0.447, 0.631], $t(7) = 1.000$, $p = .351$. A paired t test revealed the absence of any differences between the two control trials, $t(7) = 0.099$, $p = .924$.

In the test trials, individual analyses revealed that tortoises did not significantly select one food portion more than the other (Table 2). This result is confirmed by group analyses ($M = 0.516$, 95% CI

[0.449, 0.583], one-sample Wilcoxon signed-ranks test $p = .705$; Figure 4b).

Discussion

The present study represents the first attempt to investigate whether reptiles perceive visual illusions. To achieve this goal, we tested two species, *Chelonoidis carbonaria* and *Pogona vitticeps*, with one of the most popular size illusions from the human literature, the Delboeuf illusion. The findings revealed that the bearded dragons were susceptible to the illusion and appeared to perceive it in a similar manner to humans and some other species (for example, chimpanzees [Parrish & Beran, 2014] and capuchin monkeys [Parrish et al., 2015]); however, the tortoises did not

select the larger food portion when they differed in reality or when the difference was (potentially) illusory.

In control trials, 10 out of 12 bearded dragons significantly selected the larger food portion in Control A and five out of 12 in Control B. Bearded dragons' group analysis confirmed the discrimination of the larger quantity with no difference in the two control trials. When presented with the test trials, individual (seven subjects out of 12) and group analysis showed that bearded dragons selected the food portion included in the smaller plate, suggesting a human-like perception of the Delboeuf illusion. The results of Control C trials, where there were no white plates, clearly demonstrated that subjects were not basing their choice on the basis of the quantity of white visible.

Our study provides the first evidence that a reptile species, *Pogona vitticeps*, perceives a visual illusion. This indicates that, like some mammals, birds, and fish, some reptiles can interpret and alter visual input related to object size, rather than detecting visual inputs with little or no variability. The fact that bearded dragons perceive this illusion suggests the existence of assimilation and contrast phenomena in reptiles, raising the intriguing possibility that these perceptive mechanisms could be widely shared in the animal world. Also, the illusory phenomenon requires the overall perception of the array (food portion and the surrounding array). However, only few studies investigated the global and local precedence in animals. Chimpanzees and redbellied splitfin fish, like humans, display a rather robust global-to-local precedence (Fujita & Matsuzawa, 1990; Hopkins, 1997; Hopkins & Washburn, 2002; Kimchi, 1992; Navon, 1977; Truppa, Sovrano, Spinozzi, & Bisazza, 2010), and they both demonstrated perception of the Delboeuf and the Ebbinghaus illusion, respectively, as human observers. In contrast, local-to-global precedence was found in other species (e.g., capuchin monkeys: De Lillo, Spinozzi, Palumbo, & Giustino, 2011, and pigeons: Cavoto & Cook, 2001) that do not display susceptibility to the Delboeuf illusion. In this context, dogs (Miletto Petrazzini, Bisazza, & Agrillo, 2017) showed a larger interindividual variability than humans and nonsignificant trend for global precedence (Pitteri, Mongillo, Camier, & Marinelli, 2014). Given the lack of studies investigating this phenomenon, no firm conclusion can be drawn regarding the relationship between the global/local precedence and the perception of the Delboeuf illusion. What can be claimed is that the global-to-local precedence is a prerequisite but not a predictor of the illusory sensitivity. Although no study has investigated whether reptiles show global-to-local precedence, the results of our study suggest the existence of a global-to-local precedence in bearded dragons.

In contrast, this procedure proved to be less successful for studying the perception of the Delboeuf illusion in red-footed tortoises. In control trials, only two tortoises (Savina and Charles Darwin) significantly selected the larger food portion in Control A, and only one (Gerard) in Control B; in Control A, one tortoise (Ranieri) selected the smaller one instead. This performance was confirmed by group analyses that indicated no significant choice for any food portion in both types of control trials. These results suggest that, within this context, the tortoises did not maximize their food intake; therefore, we cannot draw any conclusion about their perception of the Delboeuf illusion. Why did tortoises exhibit such a poor performance? This species has been shown to optimize food intake previously and will work harder for both larger and favored food rewards, and further, they are able to retain information about stimuli associated with a greater food reward for at least

18 months (Soldati et al., 2017). It is possible that because both portion sizes used in this experiment were large (and were substantially larger than those used by Soldati et al., 2017) that either they did not need to maximize their intake or they were trying to maximize food intake, but the physical difference between the food portions presented in control trials (ratio of 0.67) was too subtle to be detected. There is currently no literature regarding the discrimination ratios in this species. If the red-footed tortoises did not discriminate a 0.67 ratio, it is likely that they would not perceive a subjective difference between the two food portions in the Delboeuf pattern. The position of the stimuli in the experimental arena could be another factor influencing their performance. Although the ramp was designed to maximize the animals' view of the entire array, perhaps this was not optimal for the tortoises.

Finally, we acknowledge our limited sample size of tortoises ($N = 8$)—although similar to the sample size used in the previous studies that investigated the spontaneous emergence of the Delboeuf illusion in other species (Miletto Petrazzini, Bisazza, & Agrillo, 2017; Parrish & Beran, 2014). Despite the limitations of a small sample, we suggest that the null results observed in tortoises could easily reflect an unidentified feature of our methodology. After determining red-footed tortoises' size discrimination abilities, one could, adopting different methodologies (e.g., training procedures or different size/position of the stimuli), investigate again their perception of the Delboeuf illusion.

The importance of reducing methodological variability is a well-known and highly debated issue in scientific community, especially in the investigation of cognitive abilities of vertebrates (Agrillo & Bisazza, 2014; Feng et al., 2017). However, as emerged in this study, the use of the same procedure and setting may prevent a reliable comparison. The use of multiple methodological approaches (e.g., free choice tests vs. operant conditioning procedure and use of food vs. two-dimensional figures as stimuli) is necessary to compare the subjective world of different species.

References

- Agrillo, C., & Bisazza, A. (2014). Spontaneous versus trained numerical abilities: A comparison between the two main tools to study numerical competence in non-human animals. *Journal of Neuroscience Methods*, 234, 82–91. <http://dx.doi.org/10.1016/j.jneumeth.2014.04.027>
- Agrillo, C., Miletto Petrazzini, M. E., & Bisazza, A. (2016). Brightness illusion in the guppy (*Poecilia reticulata*). *Journal of Comparative Psychology*, 130, 55–61. <http://dx.doi.org/10.1037/com0000020>
- Agrillo, C., Parrish, A. E., & Beran, M. J. (2014a). Do rhesus monkeys (*Macaca mulatta*) perceive the Zöllner illusion? *Psychonomic Bulletin and Review*, 21, 986–994. <http://dx.doi.org/10.3758/s13423-013-0573-2>
- Agrillo, C., Parrish, A. E., & Beran, M. J. (2014b). Do primates see the solitary illusion differently? A comparative assessment of humans (*Homo sapiens*), chimpanzees (*Pan troglodytes*), rhesus monkeys (*Macaca mulatta*), and capuchin monkeys (*Cebus apella*). *Journal of Comparative Psychology*, 128, 402–413. <http://dx.doi.org/10.1037/a0037499>
- Bánszegi, O., Urrutia, A., Szenczi, P., & Hudson, R. (2016). More or less: Spontaneous quantity discrimination in the domestic cat. *Animal Cognition*, 19, 879–888. <http://dx.doi.org/10.1007/s10071-016-0985-2>
- Bayne, K. L., & Davis, R. (1983). Susceptibility of rhesus monkeys (*Macaca mulatta*) to the Ponzo illusion. *Bulletin of the Psychonomic Society*, 21, 476–478. <http://dx.doi.org/10.3758/BF03330013>
- Byosiére, S. E., Feng, L. C., Woodhead, J. K., Rutter, N. J., Chouinard, P. A., Howell, T. J., & Bennett, P. C. (2017). Visual perception in domestic dogs:

- Susceptibility to the Ebbinghaus-Titchener and Delboeuf illusions. *Animal Cognition*, 20, 435–448. <http://dx.doi.org/10.1007/s10071-016-1067-1>
- Cavoto, K. K., & Cook, R. G. (2001). Cognitive precedence for local information in hierarchical stimulus processing by pigeons. *Journal of Experimental Psychology: Animal Behavior Processes*, 27, 3–16. <http://dx.doi.org/10.1037/0097-7403.27.1.3>
- Davis, B., Payne, C. R., & Bui, M. (2016). Making small food units seem regular: How larger table size reduces calories to be consumed. *Journal of the Association of Consumer Research*, 1, 115–124. <http://dx.doi.org/10.1086/684527>
- De Lillo, C., Spinozzi, G., Palumbo, M., & Giustino, G. (2011). Attention allocation modulates the processing of hierarchical visual patterns: A comparative analysis of capuchin monkeys (*Cebus apella*) and humans. *Journal of Experimental Psychology: Animal Behavior Processes*, 37, 341–352. <http://dx.doi.org/10.1037/a0022989>
- Feng, L. C., Chouinard, P. A., Howell, T. J., & Bennett, P. C. (2017). Why do animals differ in their susceptibility to geometrical illusions? *Psychonomic Bulletin and Review*, 24, 262–276. <http://dx.doi.org/10.3758/s13423-016-1133-3>
- Fujita, K. (1997). Perception of the Ponzo illusion by rhesus monkeys, chimpanzees, and humans: Similarity and difference in the three primate species. *Perception and Psychophysics*, 59, 284–292. <http://dx.doi.org/10.3758/BF03211896>
- Fujita, K., & Matsuzawa, T. (1990). Delayed figure reconstruction by a chimpanzee (*Pan troglodytes*) and humans (*Homo sapiens*). *Journal of Comparative Psychology*, 104, 345–351. <http://dx.doi.org/10.1037/0735-7036.104.4.345>
- Fuss, T., Bleckmann, H., & Schlüssel, V. (2014). The brain creates illusions not just for us: Sharks (*Chiloscyllium griseum*) can “see the magic” as well. *Frontiers in Neural Circuits*, 8, 24. <http://dx.doi.org/10.3389/fncir.2014.00024>
- Fuss, T., & Schlüssel, V. (2017). The Ebbinghaus illusion in the gray bamboo shark (*Chiloscyllium griseum*) in comparison to the teleost damselfish (*Chromis chromis*). *Zoology*, 123, 16–29. <http://dx.doi.org/10.1016/j.zool.2017.05.006>
- Gori, S., Agrillo, C., Dadda, M., & Bisazza, A. (2015). Do fish perceive illusory motion? *Scientific Reports*, 4, 6443. <http://dx.doi.org/10.1038/srep06443>
- Hopkins, W. D. (1997). Hemispheric specialization for local and global processing of hierarchical visual stimuli in chimpanzees (*Pan troglodytes*). *Neuropsychologia*, 35, 343–348. [http://dx.doi.org/10.1016/S0028-3932\(96\)00089-9](http://dx.doi.org/10.1016/S0028-3932(96)00089-9)
- Hopkins, W. D., & Washburn, D. A. (2002). Matching visual stimuli on the basis of global and local features by chimpanzees (*Pan troglodytes*) and rhesus monkeys (*Macaca mulatta*). *Animal Cognition*, 5, 27–31. <http://dx.doi.org/10.1007/s10071-001-0121-8>
- Horridge, G. A., Zhang, S. W., & O’Carroll, D. (1992). Insect perception of illusory contours. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences*, 337, 59–64. <http://dx.doi.org/10.1098/rstb.1992.0083>
- Howard, S. R., Avarguès-Weber, A., Garcia, J. E., Stuart-Fox, D., & Dyer, A. G. (2017). Perception of contextual size illusions by honeybees in restricted and unrestricted viewing conditions. *Proceedings of the Royal Society B*, 284, 20172278. <http://dx.doi.org/10.1098/rspb.2017.2278>
- Keep, B., Zulch, H. E., & Wilkinson, A. (2018). Truth is in the eye of the beholder: Perception of the Müller-Lyer illusion in dogs. *Learning and Behavior*, 46, 501–512. <http://dx.doi.org/10.3758/s13420-018-0344-z>
- Kelley, L. A., & Kelley, J. L. (2014). Animal visual illusion and confusion: The importance of a perceptual perspective. *Behavioral Ecology*, 25, 450–463. <http://dx.doi.org/10.1093/beheco/art118>
- Kimchi, R. (1992). Primacy of wholistic processing and global/local paradigm: A critical review. *Psychological Bulletin*, 112, 24–38. <http://dx.doi.org/10.1037/0033-2909.112.1.24>
- King, D. L. (1988). Assimilation is due to one perceived whole and contrast is due to two perceived wholes. *New Ideas in Psychology*, 6, 277–288. [http://dx.doi.org/10.1016/0732-118X\(88\)90039-6](http://dx.doi.org/10.1016/0732-118X(88)90039-6)
- Kis, A., Huber, L., & Wilkinson, A. (2015). Social learning by imitation in a reptile (*Pogona vitticeps*). *Animal Cognition*, 18, 325–331. <http://dx.doi.org/10.1007/s10071-014-0803-7>
- Lazareva, O. F., Shimizu, T., & Wasserman, E. A. (Eds.). (2012). *How animals see the world: Comparative behavior, biology, and evolution of vision*. New York, NY: Oxford University Press. <http://dx.doi.org/10.1093/acprof:oso/9780195334654.001.0001>
- Lucon-Xiccato, T., Miletto Petrazzini, M. E., Agrillo, C., & Bisazza, A. (2015). Guppies discriminate between two quantities of food items but prioritize item size over total amount. *Animal Behaviour*, 107, 183–191. <http://dx.doi.org/10.1016/j.anbehav.2015.06.019>
- Matsubara, S., Deeming, D. C., & Wilkinson, A. (2017). Cold-blooded cognition: New directions in reptile cognition. *Current Opinion in Behavioral Sciences*, 16, 126–130. <http://dx.doi.org/10.1016/j.cobeha.2017.06.006>
- Miletto Petrazzini, M. E., Bisazza, A., & Agrillo, C. (2017). Do domestic dogs (*Canis lupus familiaris*) perceive the Delboeuf illusion? *Animal Cognition*, 20, 427–434. <http://dx.doi.org/10.1007/s10071-016-1066-2>
- Miletto Petrazzini, M. E., Fraccaroli, I., Gariboldi, F., Agrillo, C., Bisazza, A., Bertolucci, C., & Foà, A. (2017). Quantitative abilities in a reptile (*Podarcis sicula*). *Biology Letters*, 13, 20160899. <http://dx.doi.org/10.1098/rsbl.2016.0899>
- Miletto Petrazzini, M. E., & Wynne, C. D. (2016). What counts for dogs (*Canis lupus familiaris*) in a quantity discrimination task? *Behavioural Processes*, 122, 90–97. <http://dx.doi.org/10.1016/j.beproc.2015.11.013>
- Moszuti, S. A., Wilkinson, A., & Burman, O. H. (2017). Response to novelty as an indicator of reptile welfare. *Applied Animal Behaviour Science*, 193, 98–103. <http://dx.doi.org/10.1016/j.applanim.2017.03.018>
- Mueller-Paul, J., Wilkinson, A., Aust, U., Steurer, M., Hall, G., & Huber, L. (2014). Touchscreen performance and knowledge transfer in the red-footed tortoise (*Chelonoidis carbonaria*). *Behavioural Processes*, 106, 187–192. <http://dx.doi.org/10.1016/j.beproc.2014.06.003>
- Murayama, T., Usui, A., Takeda, E., Kato, K., & Maejima, K. (2012). Relative size discrimination and perception of the Ebbinghaus illusion in a bottlenose dolphin (*Tursiops truncatus*). *Aquatic Mammals*, 38, 333–342. <http://dx.doi.org/10.1578/AM.38.4.2012.333>
- Nakamura, N., Watanabe, S., & Fujita, K. (2008). Pigeons perceive the Ebbinghaus-Titchener circles as an assimilation illusion. *Journal of Experimental Psychology: Animal Behavior Processes*, 34, 375–387. <http://dx.doi.org/10.1037/0097-7403.34.3.375>
- Nakamura, N., Watanabe, S., & Fujita, K. (2014). A reversed Ebbinghaus-Titchener illusion in bantams (*Gallus gallus domesticus*). *Animal Cognition*, 17, 471–481. <http://dx.doi.org/10.1007/s10071-013-0679-y>
- Navon, D. (1977). Forest before trees: The precedence of global features in visual perception. *Cognitive Psychology*, 9, 353–383. [http://dx.doi.org/10.1016/0010-0285\(77\)90012-3](http://dx.doi.org/10.1016/0010-0285(77)90012-3)
- Parrish, A. E., Agrillo, C., Perdue, B. M., & Beran, M. J. (2016). The elusive illusion: Do children (*Homo sapiens*) and capuchin monkeys (*Cebus apella*) see the Solitaire illusion? *Journal of Experimental Child Psychology*, 142, 83–95. <http://dx.doi.org/10.1016/j.jecp.2015.09.021>
- Parrish, A. E., & Beran, M. J. (2014). When less is more: Like humans, chimpanzees (*Pan troglodytes*) misperceive food amounts based on plate size. *Animal Cognition*, 17, 427–434. <http://dx.doi.org/10.1007/s10071-013-0674-3>
- Parrish, A. E., Brosnan, S. F., & Beran, M. J. (2015). Do you see what I see? A comparative investigation of the Delboeuf illusion in humans (*Homo sapiens*), rhesus monkeys (*Macaca mulatta*), and capuchin monkeys (*Cebus apella*). *Journal of Experimental Psychology: Animal Learning and Cognition*, 41, 395–405. <http://dx.doi.org/10.1037/xan0000078>

- Parron, C., & Fagot, J. (2007). Comparison of grouping abilities in humans (*Homo sapiens*) and baboons (*Papio papio*) with the Ebbinghaus illusion. *Journal of Comparative Psychology*, 121, 405–411. <http://dx.doi.org/10.1037/0735-7036.121.4.405>
- Pepperberg, I. M., & Nakayama, K. (2016). Robust representation of shape in a Grey parrot (*Psittacus erithacus*). *Cognition*, 153, 146–160. <http://dx.doi.org/10.1016/j.cognition.2016.04.014>
- Pepperberg, I. M., Vicinay, J., & Cavanagh, P. (2008). Processing of the Müller-Lyer illusion by a Grey parrot (*Psittacus erithacus*). *Perception*, 37, 765–781. <http://dx.doi.org/10.1068/p5898>
- Piaget, J. (1957). *Les Mécanismes Perceptifs*. Paris, France: Presses Universitaires de France.
- Pitteri, E., Mongillo, P., Carnier, P., & Marinelli, L. (2014). Hierarchical stimulus processing by dogs (*Canis familiaris*). *Animal Cognition*, 17, 869–877. <http://dx.doi.org/10.1007/s10071-013-0720-1>
- Pressey, A. W. (1977). Measuring the Titchener circles and Delboeuf illusions with the method of adjustment. *Bulletin of the Psychonomic Society*, 10, 118–120. <http://dx.doi.org/10.3758/BF03329298>
- Rosa Salva, O., Rugani, R., Cavazzana, A., Regolin, L., & Vallortigara, G. (2013). Perception of the Ebbinghaus illusion in four-day-old domestic chicks (*Gallus gallus*). *Animal Cognition*, 16, 895–906. <http://dx.doi.org/10.1007/s10071-013-0622-2>
- Santacà, M., Regaiolli, B., Miletto Petrazzini, M. E., Spiezio, C., & Agrillo, C. (2017). Preliminary study to investigate the Delboeuf illusion in ring-tailed lemurs (*Lemur catta*): Methodological challenges. *Animal Behavior and Cognition*, 4, 365–377. <http://dx.doi.org/10.26451/abc.04.03.13.2017>
- Siviter, H., Deeming, C. D., Rosenberger, J., Burman, O. H. P., Moszuti, S. A., & Wilkinson, A. (2017). The impact of egg incubation temperature on the personality of oviparous reptiles. *Animal Cognition*, 20, 109–116. <http://dx.doi.org/10.1007/s10071-016-1030-1>
- Siviter, H., Deeming, D. C., van Giezen, M. F. T., & Wilkinson, A. (2018). Incubation environment impacts the social cognition of adult lizards. *Royal Society Open Science*, 22, 170742. <http://dx.doi.org/10.1098/rsos.170742>
- Soldati, F., Burman, O. H. P., John, E. A., Pike, T. W., & Wilkinson, A. (2017). Long-term memory of relative reward values. *Biology Letters*, 13, 20160853. <http://dx.doi.org/10.1098/rsbl.2016.0853>
- Sovrano, V. A., Albertazzi, L., & Rosa Salva, O. (2015). The Ebbinghaus illusion in a fish (*Xenotoca eiseni*). *Animal Cognition*, 18, 533–542. <http://dx.doi.org/10.1007/s10071-014-0821-5>
- Sovrano, V. A., & Bisazza, A. (2009). Perception of subjective contours in fish. *Perception*, 38, 579–590. <http://dx.doi.org/10.1068/p6121>
- Sovrano, V. A., da Pos, O., & Albertazzi, L. (2016). The Müller-Lyer illusion in the teleost fish *Xenotoca eiseni*. *Animal Cognition*, 19, 123–132. <http://dx.doi.org/10.1007/s10071-015-0917-6>
- Suganuma, E., Pessoa, V. F., Monge-Fuentes, V., Castro, B. M., & Tavares, M. C. H. (2007). Perception of the Müller-Lyer illusion in capuchin monkeys (*Cebus apella*). *Behavioural Brain Research*, 182, 67–72. <http://dx.doi.org/10.1016/j.bbr.2007.05.014>
- Truppa, V., Sovrano, V. A., Spinozzi, G., & Bisazza, A. (2010). Processing of visual hierarchical stimuli by fish (*Xenotoca eiseni*). *Behavioural Brain Research*, 207, 51–60. <http://dx.doi.org/10.1016/j.bbr.2009.09.039>
- van Ittersum, K., & Wansink, B. (2007). Do children really prefer large portions? Visual illusions bias their estimates and intake. *Journal of the American Dietetic Association*, 107, 1107–1110. <http://dx.doi.org/10.1016/j.jada.2007.05.020>
- Wansink, B., & Cheney, M. M. (2005). Super Bowls: Serving bowl size and food consumption. *Journal of the American Medical Association*, 293, 1727–1728. <http://dx.doi.org/10.1001/jama.293.14.1727>
- Watanabe, S., Nakamura, N., & Fujita, K. (2011). Pigeons perceive a reversed Zöllner illusion. *Cognition*, 119, 137–141. <http://dx.doi.org/10.1016/j.cognition.2010.10.020>
- Watanabe, S., Nakamura, N., & Fujita, K. (2013). Bantams (*Gallus gallus domesticus*) also perceive a reversed Zöllner illusion. *Animal Cognition*, 16, 109–115. <http://dx.doi.org/10.1007/s10071-012-0556-0>
- Wilkinson, A., & Glass, E. (2018). Cold-blooded Cognition: How to get a tortoise out of its shell. In N. Bueno-Guerra & F. Amici (Eds.), *Field and laboratory methods in animal cognition* (pp. 401–422). United Kingdom: Cambridge University Press.
- Wilkinson, A., Mueller-Paul, J., & Huber, L. (2013). Picture-object recognition in the tortoise *Chelonoidis carbonaria*. *Animal Cognition*, 16, 99–107. <http://dx.doi.org/10.1007/s10071-012-0555-1>
- Wilkinson, A., Sebanz, N., Mandl, I., & Huber, L. (2011). No evidence of contagious yawning in the red-footed tortoise *Geochelone carbonaria*. *Current Zoology*, 57, 477–484. <http://dx.doi.org/10.1093/czoolo/57.4.477>
- Wyzisk, K., & Neumeier, C. (2007). Perception of illusory surfaces and contours in goldfish. *Visual Neuroscience*, 24, 291–298. <http://dx.doi.org/10.1017/S095252380707023X>

Received June 11, 2018

Revision received January 31, 2019

Accepted February 1, 2019 ■