

Predicting Longitudinal Patterns of Functional Deficits in Children With Traumatic Brain Injury

Taryn B. Fay and Keith Owen Yeates

The Ohio State University and Nationwide Children's Hospital

Shari L. Wade

University of Cincinnati and Cincinnati Children's
Hospital Medical Center

Dennis Drotar

Case Western Reserve University and Rainbow Babies &
Children's Hospital

Terry Stancin

Case Western Reserve University and
MetroHealth Medical Center

H. Gerry Taylor

Case Western Reserve University and Rainbow Babies & Children's Hospital

Longitudinal patterns of functional deficits were investigated in 37 children with severe traumatic brain injury (TBI), 40 children with moderate TBI, and 44 children with orthopedic injuries. They were from 6 to 12 years of age when injured. Their neuropsychological, behavioral, adaptive, and academic functioning were assessed at 6 months, 12 months, and 3–5 years postinjury. Functional deficits (<10th percentile for age) were identified within each outcome domain at each occasion. Children were classified into 4 a priori longitudinal patterns of outcomes within domains (i.e., no deficits, improvement, deterioration, persistent deficits). In multinomial logistic regression analyses, severe TBI predicted an increased likelihood of persistent deficits in all outcome domains, as well as deterioration in behavioral functioning and improvement in neuropsychological, adaptive, and academic functioning. Severe TBI also predicted a greater total number of functional deficits across domains at each occasion. However, many children with severe TBI showed no deficits from 6 months to 4 years postinjury in 1 or more outcome domains. The findings help clarify the course of recovery for individual children following TBI.

Keywords: traumatic brain injury, children, functional deficits, outcomes

Childhood traumatic brain injury (TBI) is associated with a variety of cognitive, behavioral, adaptive, and academic performance deficits in school-age children. Numerous studies have established persistent

neuropsychological sequelae; higher incidences of postinjury psychiatric, emotional, and behavioral problems; difficulties in adaptive functioning; and academic performance deficits after moderate to severe TBI in children (V. A. Anderson, Catroppa, Haritou, Morse, & Rosenfeld, 2005; Fay et al., 1994; Fletcher, Ewing-Cobbs, Miner, Levin, & Eisenberg, 1990; Jaffe, Polissar, Fay, & Liao, 1995; Kinsella et al., 1995; Max et al., 1997; Schwartz et al., 2003; Taylor et al., 1999, 2002; Yeates et al., 2001, 2002; Yeates & Taylor, 2006).

As a group, children with TBI show distinct longitudinal patterns of change in cognitive, academic, behavioral, and adaptive domains. Children with moderate to severe TBI generally show significant improvement in their neuropsychological functioning during the first year postinjury, yet recovery often begins to plateau after the first year postinjury, with negligible rates of change appearing during the following 2 years postinjury (Chadwick, Rutter, Brown, Shaffer, & Traum, 1981; Jaffe et al., 1995). Persistent neuropsychological sequelae of childhood TBI continuing past 1 year postinjury have also been documented (Chadwick et al., 1981; Jaffe et al., 1995; Kinsella et al., 1995; Yeates et al., 2002).

Because recovery is a developmental process that occurs over time, new impairments may also emerge (V. A. Anderson et al., 1997; Dennis, Wilkinson, Koski, & Humphreys, 1995). Long-term follow-up of children with severe TBI suggests that behavioral, adaptive, and academic deficits often fail to resolve over time, despite partial recovery in cognitive functions during the first year (Anderson et al., 2005; Jaffe et al., 1995; Kinsella, Ong, Murtaugh, Prior, & Sawyer, 1999; Perrot, Taylor, & Montes, 1991; Schwartz et al., 2003; Taylor et al., 1999, 2002; Yeates et al., 2002). Many studies have found evidence of deterioration in behavioral adjustment across time and persistent postinjury behavior problems in

Taryn B. Fay, Departments of Psychology and Pediatrics, The Ohio State University, and The Research Institute at Nationwide Children's Hospital, Columbus, Ohio; Keith Owen Yeates, Department of Pediatrics, The Ohio State University, and The Research Institute at Nationwide Children's Hospital, Columbus, Ohio; Shari L. Wade, College of Medicine, University of Cincinnati, and Department of Physical Medicine and Rehabilitation, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; Dennis Drotar and H. Gerry Taylor, Department of Pediatrics, Case Western Reserve University, and Department of Pediatrics, Rainbow Babies & Children's Hospital, Cleveland, Ohio; Terry Stancin, Department of Pediatrics, Case Western Reserve University, and Department of Pediatrics, MetroHealth Medical Center, Cleveland, Ohio.

This work was supported by National Institute of Neurological Disorders and Stroke Grant NS36335, National Institutes of Health, and by Grant MCJ-390611, Maternal and Child Health Bureau (Title V, Social Security Act, Health Resources and Services Administration, Department of Health and Human Services). We wish to acknowledge the contributions of Elizabeth Shaver, Barbara Shapero, Madeline Polonia, Marla Kemmerer, Matt Diamond, and Nichole Wood in data collection and coding. We also wish to acknowledge the participation of the Children's Hospital Medical Center of Akron and the collaboration of George Thompson, G. Dean Timmons, and Dennis Weiner. Assistance with recruitment was provided by the Rainbow Pediatric Trauma Center, Rainbow Babies & Children's Hospital, Columbus Children's Hospital Trauma Program, and MetroHealth Center Trauma Registry.

Correspondence concerning this article should be addressed to Keith Owen Yeates, Department of Psychology, Nationwide Children's Hospital, 700 Children's Drive, Columbus, OH 43205. E-mail: keith.yeates@nationwidechildrens.org

children with TBI (Jaffe et al., 1995; Schwartz et al., 2003; Taylor et al., 1999, 2001). Adaptive functioning also may decline during the first year postinjury, particularly in the domain of socialization (V. A. Anderson et al., 1997; Stancin et al., 2002). Yeates et al. (2001) demonstrated that, whereas cognitive and somatic symptoms declined during the first year postinjury in children with TBI, emotional and behavioral symptoms became more common. In addition, a number of studies have demonstrated that initial improvements in academic skills may be most prominent in children with severe TBI, with persisting or worsening functional deficits emerging at later time points (Barry, Taylor, Klein, & Yeates, 1996; Ewing-Cobbs, Fletcher, Levin, Iovino, & Miner, 1998; Jaffe et al., 1995; Max et al., 1997).

Almost all previous studies on the outcomes of TBI have compared children with TBI to those with injuries not involving the head or healthy children by testing the statistical significance of mean differences between groups, rather than by examining the presence of functional deficits in individual children. Very few studies have reported the proportion of individual children who display significant functional deficits in specific outcome domains after TBI. The few studies that have explored individual functional deficits in childhood TBI have focused on behavioral and academic impairments. For example, Max et al. (1997) found that by the second year postinjury, 36% of children with moderate to severe TBI met criteria for a new psychiatric disorder. Schwartz et al. (2003) found higher rates of clinically significant behavior problems at 3 years postinjury in children with moderate to severe TBI than in children with orthopedic injuries (OI). Similarly, V. A. Anderson et al. (2005) found that the percentage of moderate to severe behavioral impairment in children with severe TBI increased from 4.8% immediately postinjury to 34.6% by 30 months postinjury. Ewing-Cobbs et al. (1998) examined school performance in children with severe TBI and found that 79% of the cohort had either repeated a grade or were receiving special assistance in school by a 2-year postinjury follow-up. In addition, Taylor et al. (2003) investigated the rates of placement in special education programs following TBI and found that half of the severe TBI group were in special education programs several years postinjury.

In the current study, we investigated longitudinal patterns of functional deficits in children with moderate to severe TBI as compared to children with OI. Our study differs from most previous research by focusing on individual children's performance across the broad array of domains that are often examined by neuropsychologists in clinical practice, including behavioral and adaptive functioning and academic and neuropsychological performance. Moreover, we did not examine the statistical significance of group differences in means between children with TBI and OI using a variable-centered approach, as past research has done almost exclusively; instead, we adopted a person-centered approach and studied patterns of functional deficits over time in individual children. Person- and variable-centered approaches are associated with complementary analytic methods that conceptualize the organization of longitudinal data differently. Both approaches contribute to the understanding of specific links between predictors and outcomes, relations between individual positions on latent dimensions, and patterns of individual development across time (Laursen & Hoff, 2006; Magnusson, 2003; Bergman & Trost, 2006; von Eye & Bogat, 2006).

For our purposes, we identified individual patterns of functional deficits across time by choosing a cutoff score. The use of a cutoff score was necessary to classify each child's performance in terms of whether or not it represented a deficit that was atypical from a normative perspective, and hence likely to be apparent in "real-world" functioning. No universal agreement exists regarding the definition of functional deficits or clinically relevant impairments. Furthermore, cutoff scores unavoidably impose an arbitrary group structure on continuous distributions (Francis et al., 2005). However, scores that fall below the 10th percentile (i.e., ≤ -1.3 standard deviations) are often considered to represent significant impairment or deficits in clinical neuropsychological practice (Golden, Purish, & Hammeke, 1991; Hannay & Lezak, 2004; Heaton, Grant, & Matthews, 1991). We thus defined a functional deficit in this manner for the neuropsychological, academic, and adaptive domains, and in a similar manner for the behavior domain (i.e., $\geq +1.3$ standard deviations). We then classified children into four *a priori* longitudinal patterns of outcomes within each domain: no deficits, persistent deficits, improvement (initial deficit followed by no deficit), and deterioration (no initial deficit followed by later deficit).

Based on previous research, we hypothesized that injury severity would be a predictor of persistent individual deficits in all domains (V. A. Anderson et al., 1997, 2005; Fay et al., 1994; Jaffe et al., 1995; Taylor et al., 1999). Because previous literature has shown declines in behavioral functioning, we also hypothesized that children with moderate to severe TBI would be more likely to demonstrate deterioration in behavioral functioning across time than children in the OI group (V. A. Anderson et al., 2005; Max et al., 1997; Taylor et al., 2002; Yeates et al., 2001). Finally, in accordance with previous research (Jaffe et al., 1995; Taylor et al., 2002), we expected that improvements in neuropsychological and academic functioning would be more common among children with severe TBI than among children with OI.

Method

Study Design and Procedure

The study used a concurrent cohort-prospective design in which children 6 to 12 years of age were followed for approximately 3 to 5 years postinjury. The total sample included 109 children with TBI and 80 children with OI recruited from admissions to four hospitals in north-central Ohio between 1992 and 1995. Children who were admitted to the hospital with a moderate to severe TBI or OI and who met criteria for inclusion were contacted prospectively by the study coordinator at each site. An initial assessment was completed on average 3 weeks postinjury (mean interval = 3.1 weeks, $SD = 1.5$). Children with TBI had to demonstrate recovery from posttraumatic amnesia prior to any testing; only 1 child was excluded because of persistent posttraumatic amnesia that did not resolve within the necessary time frame, and that child was not testable because of a persistent vegetative state. Estimates of childhood preinjury behavioral and adaptive functioning and preinjury family functioning were obtained for both groups, based on retrospective parent ratings collected during or shortly after hospitalization. Child and family outcomes were assessed 6 and 12 months after the initial evaluation.

and again at an extended follow-up on average 4 years postinjury ($M = 4.16$ years, $SD = 0.91$). Data included in this study were obtained in compliance with the appropriate Institutional Review Boards.

Research Participants

Recruitment occurred after informed consent was obtained from the child's parent or guardian. Participants were required to (a) be between 6 and 12 years of age at the time of the injury, (b) reside in an English-speaking household, (c) be hospitalized for at least 1 night's duration for either TBI or OI, and (d) have no evidence of child abuse or previous neurological disorder. Children with TBI whose injuries did not fall into the general category of closed head injuries were excluded (e.g., brain injury due to near drowning or toxic exposure, projectile wounds, or stroke).

As reported previously (Taylor et al., 2002; Yeates et al., 2002), children with TBI were classified into two severity groups. Severe TBI was defined on the basis of the lowest Glasgow Coma Scale (GCS) score of 8 or less (Teasdale & Jennett, 1974). Moderate TBI was defined in terms of a GCS score of 9–12 or a score greater than 12 accompanied by complications (i.e., skull fracture, intracranial mass lesion or contusion, diffuse cerebral swelling, post-traumatic neurological abnormality, or documented loss of consciousness greater than 15 min). Although many children in the TBI group had accompanying orthopedic injuries, the OI group was limited to children without any symptoms suggestive of a possible central nervous system insult (e.g., symptoms of concussion). Children with OI were chosen as a comparison group to control for the effects of hospitalization as well as for premorbid characteristics that could increase the risk of nonintentional injury. The severity of traumatic injuries that were sustained by children in both groups was assessed using the Injury Severity Scale (ISS; Mayer, Matlack, Johnson, & Walker, 1980; Taylor et al., 1999). The ISS is an anatomical scoring system that provides an overall score for patients with multiple injuries. Each injury is assigned an Abbreviated Injury Scale (AIS) score and is allocated to one of six body regions, including the head. Only the highest AIS score for

each body region is used. The scores for the three most severely injured body regions are squared and summed to produce the ISS score.

For the purposes of the current analysis, we included only children who completed all follow-up assessments (6 months, 1 year, and 4 years postinjury). No significant differences were observed in age, gender, race, family composition, or severity of injury when comparing children with TBI who returned for follow-up testing with those who did not. However, families of children with TBI who did not return demonstrated a lower socioeconomic status. Among children with OI, no significant differences in demographic characteristics were found between those who returned for follow-up testing and those who did not. The final sample consisted of 37 children with severe TBI, 40 with moderate TBI, and 44 children with OI.

Table 1 summarizes the demographic characteristics of the three groups, which did not differ in age at injury, gender, family composition, socioeconomic status, or time from injury to the 4-year follow-up testing. Children from the OI group were more likely to be of African American ethnicity, $\chi^2(1, N = 121) = 5.97$. In addition, and as expected, the groups differed significantly in duration of hospitalization ($p < .05$) and ISS ($p < .01$), with the severe TBI group demonstrating the longest hospitalization stay and most severe injuries. The moderate TBI group had a somewhat higher mean GCS score than might be expected, but included children who in some studies would be classified as complicated mild TBI (i.e., GCS 13–15 but with extended loss of consciousness or abnormal neuroimaging).

Measures

Family functioning. Family measures included parent ratings of premorbid family functioning and non-injury-related environmental stress, obtained at the baseline assessment, as well as a composite measure of socioeconomic status. The 12-item General Functioning Scale from the McMaster Family Assessment Device was used as a summary measure of family functioning (Miller, Bishop, Epstein, & Keitner, 1985; Stevenson-Hinde & Akister, 1995). Environmental stress unrelated to the injury was measured

Table 1
Group Characteristics

Characteristic	Group		
	Orthopedic injury ($n = 44$)	Moderate TBI ($n = 40$)	Severe TBI ($n = 37$)
Mean (SD) age at injury (years)	9.47 (1.90)	9.49 (1.86)	9.77 (2.18)
Mean (SD) lowest Glasgow Coma Scale score*	n/a	14.08 (1.74)	4.70 (1.91)
Mean (SD) duration of unconsciousness (days)*	n/a	0.18 (.59)	5.11 (5.85)
Mean (SD) duration of hospitalization (days)*	15.09 (14.47)	6.80 (7.28)	13.16 (9.60)
Mean (SD) Injury Severity Scale score*	7.27 (2.77)	12.33 (5.68)	19.97 (12.09)
Sex (% boys)	61	68	81
Ethnicity (% Caucasian)*	57	75	81
Mean (SD) socioeconomic composite index	−0.01 (.90)	−0.23 (.88)	−0.15 (.87)
Family composition (% two-parent families)	57	70	68
Mean (SD) time from injury to 4-year follow-up (years)	4.18 (0.94)	4.25 (0.84)	4.04 (0.94)

Note. TBI = traumatic brain injury.

* Group differences significant, $p < .05$.

using a composite derived from the Life Stressors and Social Resources Inventory—Adult Form (LISRES–A; Moos & Moos, 1994). The composite was an average of *T* scores for all stress subscales aside from those specific to the child and to negative life events. Potential effects of the original injury on the latter two subscales justified their exclusion. A socioeconomic composite index was computed to provide a distal measure of the family environment (Yeates & Taylor, 1997). It was computed by averaging sample *z* scores for the Duncan Occupational Status Index (Stevens & Featherman, 1981), annual family income as coded on the LISRES–A, and years of maternal education as measured at the baseline assessment.

Neuropsychological functioning. Children were administered a set of neuropsychological tests (see Yeates et al., 2002, for details). The tests included a short form of the Wechsler Intelligence Scale for Children—Third Edition (Wechsler, 1991), which provided prorated verbal and performance IQ scores; Recalling Sentences subtest of the Clinical Evaluation of Language Fundamentals—Revised (Semel, Wiig, Secord, & Sabers, 1987); Developmental Test of Visual–Motor Integration (Beery, 1989); Contingency Naming Test (Taylor, Hack, & Klein, 1998; P. Anderson, Anderson, Northam, & Taylor, 2000); Boston Naming Test (Yeates, 1994); Controlled Oral Word Association Test (Spreen & Strauss, 1991); a prepublication version of the California Verbal Learning Test (Delis, Kramer, Kaplan, & Ober, 1986); Continuous Performance Test—3 (Lindgren & Lyons, 1984); Subtests 2, 4, and 9 of the Underlining Test (Rourke & Orr, 1977); and Grooved Pegboard Test (Knights & Norwood, 1980). All tests were completed at each assessment, with the exception of the Grooved Pegboard, which was administered only at the 6- and 12-month assessments.

A poor neuropsychological outcome was defined on the basis of the presence of three or more functional deficits (i.e., age-based standardized scores below the 10th percentile) at each assessment. Age-corrected standard scores were used in coding tests with published norms. For tests without published norms, raw scores were transformed to age- and gender-corrected *z* scores prior to analysis (Taylor & Schatschneider, 1992). Transformations were based on the performances of the OI group at each assessment. Eleven neuropsychological tests were administered in total; the cutoff of three or more functional deficits minimized the likelihood of false positives when identifying a poor neuropsychological outcome at each assessment (Ingraham & Aiken, 1996).

Behavioral functioning. Behavioral functioning was assessed by having parents complete the Child Behavior Checklist (Achenbach, 1991), a widely used rating scale with good reliability and validity. A functional deficit in behavioral adjustment was defined as a Total Behavior Problem *T* score above the 90th percentile.

Adaptive functioning. Adaptive functioning was assessed using the Vineland Adaptive Behavior Scales, which is a semistructured parent interview (Sparrow, Balla, & Cicchetti, 1984). A functional deficit in adaptive behavior was defined as one or more age-based standard scores below the 10th percentile on the Communication, Daily Living Skills, or Socialization scales.

Academic outcome. Children completed the Word Identification, Writing Sample, and Calculation subtests of the Woodcock–Johnson Test of Achievement—Revised (Woodcock & Mather, 1989). A functional deficit in academic skills was defined as one or more age-based standard scores below the 10th percentile.

Data Analysis

Each individual child's outcomes at each occasion within each domain (i.e., neuropsychological, behavioral, adaptive, and academic) were coded into dichotomous dependent variables (i.e., functional deficit vs. no deficit) using the criteria described above. We then examined patterns of outcomes in each domain across time. Children were classified into four a priori longitudinal patterns of outcomes within domains: no deficit at any occasion, persistent deficits at all occasions, improvement (i.e., deficit at 6 months followed by no deficit at 1 and 4 years or deficits at 6 months and 1 year followed by no deficit at 4 years), and deterioration (i.e., no deficit at 6 months followed by deficits at both 1 and 4 years postinjury or no deficits at 6 months or 1 year followed by deficit at 4 years postinjury). The classification represents a person-centered approach to outcome assessment in high-risk groups (Connell, Dishion, & Deater-Deckard, 2006; Laursen & Hoff, 2006; Masten, 2001), focusing on patterns of outcome suggested by previous longitudinal studies of children with TBI (Taylor, 2004).

Multinomial logistic regression was used to compare the TBI and OI groups in terms of the frequency of different patterns of postinjury functional deficits for each of the four types of outcomes. The analyses included two dummy variables that represented contrasts between the severe TBI and OI groups and between the moderate TBI and OI groups. Predictors used as covariates included ethnicity (White vs. non-White), socioeconomic status (socioeconomic composite index), preinjury family functioning (McMaster Family Assessment Device General Functioning), environmental stress (LISRES–A), child preinjury child behavioral adjustment and school performance (Child Behavior Checklist Total Problems *T* score and School Competence *T* score), and preinjury child adaptive functioning (baseline Vineland Adaptive Behavior Scales Adaptive Behavior Composite standard score). Preinjury child status was thus assessed using multiple indicators. The dependent variable in each analysis was the pattern of outcome, using “no deficits” as the reference category.

To identify indices of TBI severity associated with patterns of functional deficits over the follow-up interval, we conducted separate multinomial logistic regression analyses for each functional outcome for the TBI sample only. Instead of dummy variables representing group membership, the lowest postresuscitation GCS score and duration of impaired consciousness were used as indices of severity of injury. Because the two indices are highly correlated, analyses including both indices would not converge or were characterized by singularities; therefore, separate multinomial logistic regression analyses were conducted for each measure of severity. The other predictors remained largely the same as the group comparisons, although environmental stress (LISRES–A) was eliminated from the analyses because some analyses would not converge to a solution when it was included. The reference category for each dependent variable again was “no deficits.”

A variable number of children showed patterns of functional deficits that could not be classified into the four major categories and were excluded from the respective multinomial logistic regression analysis. Seven children were excluded in the neuropsychological domain (1 severe TBI, 3 moderate TBI, 3 OI), 9 in the behavioral domain (1 severe TBI, 4 moderate TBI, 4 OI), 4 in the academic domain (1 severe TBI, 1 moderate TBI, 2 OI), and 13 in

the adaptive domain (6 severe TBI, 5 moderate, 2 OI). The children excluded from one analysis were not always the same as those excluded from any other. The total number of participants reported in Table 1 includes all cases.

To examine the comorbidity of functional deficits, we conducted ordinal regression analyses on the total number of functional deficits (range = 0–4) at each of the follow-up assessments. Similar to the multinomial regression analysis, the ordinal regression analyses were conducted with the same dummy variables representing group contrasts and the same covariates. In addition, separate ordinal regression analyses were conducted on the TBI sample only with GCS score and duration of impaired consciousness used as indices of injury severity. The ordinal regression analyses included all children.

Results

Group Differences in Functional Deficits

Table 2 presents the number and percentage of children from each group who displayed the four longitudinal patterns of outcome within each domain. In the multinomial logistic regression analyses, the addition of the severe TBI and OI group contrast contributed to improved model fit in all four outcome domains (see Table 3), all χ^2 s > 8.13, and all ps < .05. As Table 2 shows, the severe TBI group was more likely than the OI group to display persistent deficits in all four outcome domains (i.e., 17%–23% vs. 3%–7%, depending on domain). In addition, the severe TBI group was more likely than the OI group to display a pattern of deterioration in behavioral outcomes (19% vs. 5%). Notably, Table 2 also shows that between 48% and 67% of the severe TBI group showed no functional deficits in neuropsychological, behavioral, academic, or adaptive functioning over time.

The addition of the moderate TBI and OI group contrast also contributed to improved model fit for the neuropsychological and academic outcomes, χ^2 s > 7.90, p < .05 (see Table 3). The moderate TBI group was significantly more likely than the OI group to display persistent neuropsychological deficits (14% vs. 5%; see Table 2). None of the specific patterns involving adaptive or academic outcomes was significantly more common among the moderate TBI group.

Children's premorbid functioning was also a consistent predictor of outcome. Premorbid adaptive functioning significantly predicted neuropsychological, behavioral, adaptive, and academic outcomes, all χ^2 s > 8.34 and all ps < .05; premorbid behavioral functioning significantly predicted behavioral and adaptive outcomes, both χ^2 s > 13.59 and both ps < .05; and premorbid school functioning significantly predicted academic outcomes, χ^2 = 22.79 and p < .01. In general, poorer premorbid functioning predicted an increased likelihood of persistent deficits (see Table 3). Poorer premorbid functioning also predicted an increased likelihood of improvement in some domains, although poor premorbid school performance predicted an increased likelihood of deterioration in academic outcomes.

Environmental factors at baseline also predicted outcomes in multiple domains. Socioeconomic status predicted neuropsychological, behavioral, and academic outcomes, both χ^2 s > 18.78 and all ps < .01, and family environmental stress (i.e., LISRES-A) predicted academic outcomes, χ^2 = 8.59 and p < .05. In general, lower socioeconomic status was associated with increased likelihood of persistent deficits, and lower family stress predicted an increased likelihood of improvement in academic functioning (see Table 3).

Table 4 shows the number and percentage of children from each group who displayed from zero to four total deficits across do-

Table 2
Patterns of Functional Deficits

Variable	Group		
	Orthopedic injury <i>n</i> (%)	Moderate TBI <i>n</i> (%)	Severe TBI <i>n</i> (%)
Neuropsychological outcome			
No deficits	33 (81)	30 (81)	24 (67)
Improvement	2 (5)	1 (3)	4 (11)
Deterioration	4 (10)	1 (3)	2 (6)
Persistent deficits	2 (5)	5 (14)	6 (17)
Behavioral outcome			
No deficits	35 (88)	28 (78)	21 (58)
Improvement	2 (5)	1 (3)	2 (6)
Deterioration	2 (5)	2 (7)	7 (19)
Persistent deficits	1 (3)	5 (14)	6 (17)
Adaptive outcome			
No deficits	27 (64)	24 (69)	15 (48)
Improvement	3 (7)	2 (6)	2 (7)
Deterioration	10 (24)	7 (20)	7 (23)
Persistent deficits	2 (5)	2 (6)	7 (23)
Academic outcome			
No deficits	34 (81)	29 (74)	21 (58)
Improvement	2 (5)	4 (10)	5 (14)
Deterioration	3 (7)	3 (8)	3 (8)
Persistent deficits	3 (7)	3 (8)	7 (19)

Note. TBI = traumatic brain injury.

Table 3

Multinomial Logistic Regression Results Showing Significant Predictors of Each Longitudinal Pattern: Total Group (Orthopedic Injury and Traumatic Brain Injury)

Outcome	Variable	B (SE)	OR (95% CI)	p
Neuropsychological outcome				
Improvement	No significant results			
Deterioration	No significant results			
Persistent deficits	Socioeconomic composite index	4.84 (1.81)	125.93 (3.63, 4374.77)	.01
	Race (1 = minority)	3.84 (1.45)	0.01 (2.70, 795.15)	.01
	School competence	−0.16 (.08)	0.86 (0.73, 0.99)	.04
	Child Behavior Checklist total	−0.15 (.07)	0.87 (0.75, 0.99)	.05
	Vineland Adaptive Behavior	−0.17 (.08)	0.84 (0.71, 0.99)	.04
	OI vs. moderate TBI	−3.55 (1.73)	0.03 (0.001, 0.85)	.04
	OI vs. severe TBI	−3.64 (1.70)	0.03 (0.001, 0.73)	.03
Behavioral outcome				
Improvement	Socioeconomic composite index	3.46 (1.62)	31.87 (1.33, 762.95)	.03
	Child Behavior Checklist total	0.25 (.09)	1.28 (1.07, 1.53)	.01
Deterioration	Child Behavior Checklist total	0.15 (.06)	1.17 (1.04, 1.32)	.01
	Vineland Adaptive Behavior	0.10 (.05)	1.10 (1.01, 1.20)	.03
	OI vs. severe TBI	−3.55 (1.23)	0.03 (0.003, 0.32)	.004
Persistent deficits	Socioeconomic composite index	4.59 (1.69)	98.04 (3.57, 2694.87)	.01
	Child Behavior Checklist total	0.37 (.11)	1.45 (1.16, 1.80)	.001
	Vineland Adaptive Behavior	0.15 (.07)	1.16 (1.02, 1.33)	.03
	OI vs. severe TBI	−6.37 (2.63)	0.002 (0.00, 0.30)	.02
Adaptive outcome				
Improvement	Child Behavior Checklist total	0.18 (.08)	1.20 (1.03, 1.40)	.02
	Vineland Adaptive Behavior	−0.21 (.10)	0.81 (0.66, 0.98)	.03
Deterioration	Race (1 = minority)	1.47 (.73)	4.36 (1.05, 18.08)	.04
	Life Stressors and Resources Inventory—Adult	0.14 (.07)	1.15 (1.01, 1.31)	.03
	School competence	−0.10 (.04)	0.90 (0.83, 0.98)	.02
Persistent deficits	Child Behavior Checklist total	0.23 (.08)	1.25 (1.07, 1.46)	.01
	OI vs. severe TBI	−3.81 (1.54)	0.02 (0.001, 0.45)	.01
Academic outcome				
Improvement	Life Stressors and Resources Inventory—Adult	−0.18 (.09)	0.83 (0.70, 0.99)	.04
	Vineland Adaptive Behavior	−0.10 (.05)	0.90 (0.82, 0.99)	.04
Deterioration	Socioeconomic composite index	3.91 (1.56)	49.89 (2.36, 1056.52)	.01
	Family functioning (FAD)	−3.96 (1.99)	0.02 (0.00, 0.96)	.05
	School competence	−0.45 (.16)	0.64 (0.47, 0.87)	.01
Persistent deficits	Socioeconomic composite index	8.37 (3.27)	4328.47 (7.19, 2604821.99)	.01
	Vineland Adaptive Behavior	−0.12 (.06)	0.89 (0.79, 0.99)	.05
	OI vs. severe TBI	−3.77 (1.68)	0.02 (0.00, 0.61)	.02

Note. OI = orthopedic injury; TBI = traumatic brain injury; FAD = McMaster Family Assessment Device.

mains at each occasion. The groups differed significantly in the total number of total functional deficits. The models that included the contrast between the severe TBI and OI groups demonstrated a significantly better overall fit than models without that contrast at all three time periods (all Wald statistics > 8.36 and all $ps < .05$). As Table 4 shows, children in the severe TBI group were more likely than those in the OI group to display three or more functional deficits at all time periods, depending on the time postinjury (i.e., 16%–25% vs. 2%–12%).

Models that included the contrasts between the moderate TBI and OI groups resulted in a significantly better overall fit than models without that contrast at only 1 year postinjury (Wald statistic = 6.92, $p < .05$; see Table 4). Children in the moderate TBI group were more likely than those in the OI group to display more total deficits (13% displayed three or more total deficits vs. 2% of the OI group).

Indices of TBI Severity as Predictors of Functional Deficits

Table 5 summarizes the results of the multinomial regression analyses using duration of unconsciousness to predict the four longitudinal patterns of outcome within each domain. Models that included duration of unconsciousness as a predictor demonstrated a significantly better overall fit than models without that predictor for neuropsychological, adaptive, and academic outcomes, $\chi^2 > 8.41$, $p < .05$. A longer duration of unconsciousness was associated with a greater likelihood of persistent neuropsychological and adaptive deficits, as well as with a greater likelihood of improvement in neuropsychological, adaptive, and academic outcomes.

Table 6 summarizes the results of the multinomial regression analyses using lowest GCS score to predict the four longitudinal patterns of outcome within each domain. Inclusion of the lowest GCS as a predictor yielded a significantly better overall fit than

Table 4
Total Number of Functional Deficits Across Domains by Group

Variable	Group		
	Orthopedic injury n (%)	Moderate TBI n (%)	Severe TBI n (%)
Total deficits at 6 months postinjury			
No deficits	31 (71)	24 (60)	17 (46)
One deficit	8 (18)	9 (23)	6 (16)
Two deficits	2 (5)	4 (10)	5 (14)
Three deficits	1 (2)	2 (5)	7 (19)
Four deficits	2 (5)	1 (3)	2 (5)
Total deficits at 1 year postinjury			
No deficits	33 (75)	22 (55)	16 (43)
One deficit	4 (9)	10 (25)	7 (19)
Two deficits	6 (14)	3 (8)	8 (22)
Three deficits	1 (2)	3 (8)	4 (11)
Four deficits	0 (0)	2 (5)	2 (5)
Total deficits at 4 years postinjury			
No deficits	26 (59)	22 (55)	14 (38)
One deficit	12 (27)	10 (25)	8 (22)
Two deficits	1 (2)	4 (10)	6 (16)
Three deficits	2 (5)	2 (5)	5 (14)
Four deficits	3 (7)	2 (5)	4 (11)

Note. TBI = traumatic brain injury.

when that predictor was excluded for behavioral, adaptive, and academic outcomes, $\chi^2 > 8.14$, $p < .05$. A lower GCS score was associated with a greater likelihood of persistent adaptive deficits and improvement in adaptive outcome, as well as with a greater likelihood of deterioration in behavioral outcomes.

When examining the TBI group only, children's premorbid functioning remained a consistent predictor of outcome. Premorbid behavioral functioning significantly predicted behavioral, adaptive, and academic outcomes, all χ^2 s > 9.13 and all $ps < .05$; premorbid school performance significantly predicted neuropsychological outcome.

Table 5
Multinomial Logistic Regression Showing Significant Predictors of Outcome: Traumatic Brain Injury Group Only
Index of Severity = Duration of Unconsciousness

Outcome	Variable	B (SE)	OR (95% CI)	p
Neuropsychological outcome				
Improvement	Duration of unconsciousness	0.34 (0.14)	1.41 (1.07, 1.85)	.01
Deterioration	No significant results			
Persistent deficits	Socioeconomic composite index	25.38 (11.66)	0.00 (12.47, 0.00)	.03
	Race (1 = minority)	34.60 (16.99)	0.00 (3.67, 0.00)	.04
	Duration of unconsciousness	2.45 (1.07)	11.55 (1.40, 95.21)	.02
Behavioral outcome				
Improvement	No significant results			
Deterioration	Socioeconomic composite index	1.71 (0.81)	5.55 (1.14, 27.03)	.03
	Child Behavior Checklist total	0.15 (0.07)	1.15 (1.01, 1.32)	.04
Persistent deficits	Socioeconomic composite index	8.30 (3.43)	4011.186 (4.84, 3325495.92)	.02
	Child Behavior Checklist total	0.48 (0.18)	1.61 (1.14, 2.28)	.01
	Race (1 = minority)	-4.94 (2.42)	0.00 (0.00, 0.83)	.01
Adaptive outcome				
Improvement	Child Behavior Checklist total	0.57 (0.22)	1.78 (1.15, 2.74)	.01
	Duration of unconsciousness	0.44 (0.17)	1.55 (1.12, 2.14)	.01
Deterioration	Child Behavior Checklist total	0.15 (0.07)	1.16 (1.02, 1.32)	.02
Persistent deficits	Socioeconomic composite index	2.88 (1.23)	17.89 (1.61, 199.14)	.02
	Child Behavior Checklist total	0.25 (0.10)	1.28 (1.05, 1.57)	.01
	Vineland Adaptive Behavior	-0.22 (0.10)	0.80 (0.66, 0.97)	.02
Academic outcome				
Improvement	Child Behavior Checklist total	0.18 (0.07)	1.20 (1.04, 1.37)	.01
	Duration of unconsciousness	0.24 (0.10)	1.27 (1.03, 1.55)	.02
Deterioration	School competence	-0.28 (0.11)	0.76 (0.61, 0.93)	.01
Persistent deficits	Socioeconomic composite index	7.57 (2.60)	1933.116 (11.75, 318073.99)	.01

Table 6

Multinomial Logistic Regression Showing Significant Predictors of Outcome: Traumatic Brain Injury Group Only
Index of Severity = Lowest Glasgow Coma Scale (GCS) Score

Outcome	Variable	B (SE)	OR (95% CI)	p
Neuropsychological outcome				
Improvement	No significant results			
Deterioration	No significant results			
Persistent deficits	Socioeconomic composite index	3.52 (1.54)	33.73 (1.66, 684.12)	.02
	Race (1 = minority)	3.46 (1.49)	31.77 (1.70, 592.79)	.02
Behavioral outcome				
Improvement	No significant results			
Deterioration	Socioeconomic composite index	2.49 (1.02)	12.09 (1.65, 88.66)	.01
	Child Behavior Checklist total	0.26 (0.11)	1.29 (1.05, 1.59)	.01
	Vineland Adaptive Behavior	0.13 (0.06)	1.14 (1.00, 1.29)	.05
	Lowest GCS score	-0.30 (0.14)	0.74 (0.57, 0.97)	.03
Persistent deficits	Socioeconomic composite index	9.36 (4.15)	11638.42 (3.40, 39879659.08)	.02
	Race (1 = minority)	-6.06 (2.96)	0.002 (0.00, 0.77)	.04
	Child Behavior Checklist total	0.60 (0.23)	1.82 (1.17, 2.84)	.01
Adaptive outcome				
Improvement	Child Behavior Checklist total	0.59 (0.19)	1.81 (1.24, 2.63)	.002
	Lowest GCS score	-0.73 (0.32)	0.48 (0.26, 0.91)	.02
Deterioration	Child Behavior Checklist total	0.19 (0.08)	1.21 (1.02, 1.43)	.02
Persistent deficits	Socioeconomic composite index	4.60 (1.83)	99.84 (2.78, 3583.95)	.01
	Child Behavior Checklist total	0.65 (0.23)	1.92 (1.23, 2.99)	.004
	Lowest GCS score	-1.17 (0.43)	0.31 (0.14, 0.72)	.01
Academic outcome				
Improvement	Child Behavior Checklist total	0.17 (0.07)	1.19 (1.04, 1.36)	.01
Deterioration	Socioeconomic composite index	12.34 (6.05)	227753.62 (1.63, 318129933.16)	.04
Persistent deficits	Socioeconomic composite index	20.35 (9.63)	0.00 (4.38, 0.00)	.04

chological, behavioral, and academic functioning, all χ^2 s > 10.91 and all ps < .05 (see Tables 5 and 6). More behavioral problems at baseline were associated with persistent deficits and deterioration in behavioral and adaptive domains and improvement in adaptive and academic domains. Poorer premorbid school performance was associated with an increased likelihood of deterioration in academic functioning. Environmental variables were less consistently associated with outcomes in the TBI group, but lower socioeconomic status predicted persistent deficits in all outcome domains, all χ^2 s > 9.64 and all ps < .05, as well as deterioration in behavioral and academic functioning (see Tables 5 and 6).

Duration of unconsciousness and lowest GCS score were also significant predictors of the number of functional deficits at each follow-up occasion. Models that included duration of unconsciousness and lowest GCS score as predictors of the total number of deficits at each occasion demonstrated a significantly better overall fit than models without these predictors, such that greater injury severity was consistently associated with a higher total number of functional deficits (all Wald statistics > 4.59 and all ps < .05).

Discussion

By focusing on individual functional deficits, the current findings help clarify the nature of the course of recovery after pediatric TBI. The findings support our central hypothesis that children with severe TBI are more likely to demonstrate persistent functional deficits in neuropsychological, behavioral, adaptive, and academic outcomes than children with OI. On the other hand, many children with severe TBI showed no deficits from 6 months to 4 years postinjury in one or more outcome domains. The apparent absence of deficits for these children does not prove that they had no

deficits, as many may have been more adversely affected prior to the 6-month follow-up or may have sustained sequelae that were too mild to meet our criteria for functional deficits. Moreover, the severe TBI group displayed significantly more total deficits at all time periods as compared with the OI group. At 1 year postinjury, nearly 60% of the severe TBI group showed functional deficits in at least one domain, as opposed to about 25% of the OI group. Similarly, at 4 years postinjury, about 40% of the severe TBI group showed functional deficits in two or more domains, whereas less than 20% of the OI group did so.

Duration of unconsciousness and lowest GCS score also were significant predictors of patterns of deficits across time and total number of deficits. These results are consistent with our previous findings from this study, as well as with other research demonstrating that injury severity is a key predictor of outcome in childhood TBI (V. A. Anderson et al., 1997, 2005; Kinsella et al., 1999; Schwartz et al., 2003; Taylor et al., 2002; Yeates et al., 2001). We were unable to compare the two measures as predictors of outcomes directly because of their high correlation. However, duration of unconsciousness was a significant predictor of outcomes in a somewhat different set of domains (i.e., neuropsychological, adaptive, academic) than was lowest GCS score (i.e., behavioral, adaptive). Thus, neither predictor appears to be consistently superior to the other, consistent with previous research on the ability of measures of TBI severity to predict neurobehavioral and functional outcomes (Massagli, Michaud, & Rivara, 1996; McDonald et al., 1994).

As hypothesized, children in the severe TBI group not only were more likely to show persistent functional deficits across domains, but also were significantly more likely to display a pattern of

deterioration in behavioral outcomes. The latter findings suggest the delayed onset of significant behavior problems for some children with TBI. This result is consistent with our previous findings, as well as with other studies showing that behavior problems in children with severe TBI persist past 1 year postinjury, and highlights the specific incidence of delayed behavior problems in individual children following a severe TBI (Max et al., 1997; Schwartz et al., 2003; Taylor et al., 1999, 2002; Yeates & Taylor, 2006). V. A. Anderson et al. (2005) specifically demonstrated comparable findings, such that children with severe TBI showed increased behavioral impairment by 30 months postinjury.

Again as predicted, we found that improvement in neuropsychological and academic domains was more common among children with severe TBI. For instance, 11% of children with severe TBI demonstrated improvement in neuropsychological functioning over time compared with only 3% of children with moderate TBI and 5% of children with OI. This is consistent with a study by Jaffe et al. (1995) that showed improvements in academic and neuropsychological functioning during the first year postinjury, with persistent deficits continuing across 3 years.

Although not a primary focus, our analyses also showed that children's premorbid functioning helped account for functional outcomes. For instance, poorer premorbid behavioral and adaptive functioning predicted persistent neuropsychological, behavioral, and adaptive deficits. Worse premorbid behavioral functioning also predicted deterioration in behavioral and adaptive functioning. These findings are in line with previous studies on this cohort by Schwartz et al. (2003) and Taylor et al. (2002), showing that preinjury behavioral adjustment predicts the onset of behavior problems after injury. V. A. Anderson et al. (1997) reported similar findings, with preinjury adaptive functioning predicting intellectual functioning at 12 months postinjury in children with moderate and severe TBI. We also found that lower premorbid school competence was associated with persistent neuropsychological deficits and deterioration in adaptive and academic functioning. In addition, low socioeconomic status was consistently associated with persistent deficits in all domains. Taken together, these findings indicate that children with poorer premorbid functioning and less advantaged family environments are at greater risk for persistent functional deficits and deterioration in many domains of functioning.

In summary, our findings build on the existing literature by demonstrating that a significant proportion of children demonstrate functional deficits following severe TBI, and those who do so are at risk for persistent deficits, even many years after recovery. The results also indicate that some children with severe TBI demonstrate recovery in academic and neuropsychological domains, but some show the delayed onset of significant behavioral problems. Children with better premorbid functioning may have a greater chance of recovery or improvement over time, whereas those with significant preinjury problems may be more likely to display negative long-term sequelae of TBI.

At the same time, the findings also indicate that many children with moderate to severe TBI actually show few functional deficits. Indeed, one of the central findings presented here is that the TBI groups demonstrate less functional impairment than might have been expected based on existing research, which generally uses a variable-centered approach that focuses on the statistical significance of group differences in mean scores. Variable-centered

analyses may lead to an overestimation of the incidence of individual deficits following TBI because the conclusions drawn are based on groups that show considerable overlap in the distribution of scores, even in the presence of a significant difference with effect sizes of the magnitude often found in studies of TBI (i.e., .5 to 1 standard deviation). Our findings are based on individual scores rather than group means, as well as on reasonably stringent definitions of clinical impairment; they therefore provide an important and different perspective on the nature of deficits following childhood TBI.

A limitation of this study is the use of a cutoff score to classify functional deficits. Cutoff scores have inherent limitations, but are a convenient means of differentiating typical from atypical performance. Any choice of a cutoff score is somewhat arbitrary, although we chose a cutoff often considered to represent significant impairment in clinical neuropsychological practice. A more liberal cutoff (e.g., 15th percentile) would have led to the identification of more functional deficits, even in the OI group. A more conservative cutoff (e.g., 5th percentile) would have resulted in the identification of few functional deficits, even among children with TBI. In either instance, however, similar differences would have been found between the TBI and OI groups in the relative incidence of deficits.

One concern about the use of cutoff scores is that they may not result in reliable classification because changes in classification can occur with relatively trivial changes in performance. We performed follow-up analyses comparing the actual performances within outcome domains of the individuals who were classified according to their patterns of functional deficits. The analyses showed that the groups displayed significantly different patterns of mean scores in each outcome domain across time (i.e., neuropsychological, behavioral, adaptive, and academic). Moreover, the mean scores showed significantly different trends across time, consistent with the patterns of functional deficits into which they were classified (i.e., no deficits, persistent deficits, improvement, and deterioration). Thus, the classification of individual functional deficits across time was associated with corresponding group change or stability in mean scores within domains.

Alternatively, we could have calculated a reliable change index (RCI) to determine whether the difference between initial and subsequent test scores from the same examinee could be attributed to measurement error (Jacobson, Roberts, Berns, & McGlinchey, 1999; Jacobson & Truax, 1991; 20Slick, 2006). However, an RCI can be used with only two time points, whereas our study examined outcome over three time points. In addition, RCI applies to a single measure, whereas some of the outcomes in the current study were assessed on the basis of multiple indices (e.g., neuropsychological functioning). Perhaps more important, a significant RCI can occur even when neither score being compared is normatively atypical, and therefore would not necessarily have functional implications (Slick, 2006). We were interested in whether or not each child demonstrated a functional deficit at each time point, not whether the change from baseline was significant. In clinical practice, neuropsychologists scrutinize individual test scores to determine whether or not a child evidences functional impairment. This study used similar methods to examine the functional outcomes of children with TBI.

The study was also limited because environmental variables were examined only as covariates in the prediction of functional outcomes, rather than as potential moderators, as we have done in past studies (Taylor et al., 2002; Yeates et al., 2002). We attempted to assess the effects of interactions of injury severity with environmental variables in the multinomial logistic regression analyses. However, in three of the four cases, entry of the interaction terms led to Hessian singularities or problems of convergence. This is most likely a function of the multicollinearity between interaction terms, as well as the relatively small number of cases within some of the groups of longitudinal patterns. Future studies with larger sample sizes will be needed to investigate the moderating effects of environmental variables on functional outcomes after childhood TBI. Another limitation in our analysis is that we did not use advanced statistical techniques that derive patterns of outcomes across time empirically (e.g., latent transition analysis, latent class growth analysis), largely because our sample size would not necessarily support such techniques (Muthén & Muthén, 2000).

Another limitation shared by many studies of childhood TBI is that children's behavioral and adaptive functioning, as well as family functioning and environmental stress, were all assessed by parent report. In addition, the parents' ratings of premorbid functioning were unavoidably retrospective. As well, families of children with TBI who did not return for follow-up were of lower socioeconomic status than those who remained in the study. Because the outcomes examined in this study are influenced by socioeconomic status, and the TBI participants who remained in the study were of higher socioeconomic status than those who did not, our results may underestimate somewhat the proportion of functional deficits that occur after pediatric TBI. However, because we controlled for socioeconomic status in our analyses and found no other demographic differences between groups, group differences in functional deficits are unlikely to be strongly affected by this bias. A final limitation is that follow-up assessments were not completed between 1 and 4 years postinjury; more frequent follow-ups would have allowed for a more precise description and categorization of longitudinal patterns of outcome.

Despite these limitations, the current results indicate that children show significant variation in their functioning after a moderate to severe TBI, with a substantial proportion actually showing relatively few functional deficits during recovery. We found that a child's individual pattern of functional deficits after a moderate to severe TBI is likely determined by a variety of factors, including severity of injury, premorbid functioning, and environmental factors such as socioeconomic status. Notably, functional deficits were not uncommon in the severe TBI group. Approximately two of five children with severe TBI demonstrated two or more functional deficits that persisted more than 4 years postinjury, despite the provision of substantial clinical and educational services to many of them. These persistent deficits highlight the need for careful evaluation and ongoing monitoring of children with severe TBI throughout their recovery given that some of them will demonstrate multiple areas of functional impairment that persist for many years after their injuries. These deficits are likely to have a significant impact on children's general quality of life and presumably will necessitate ongoing intervention (Stancin et al., 2002; Yeates &

Taylor, 2006). On the other hand, the findings hold out hope that children with severe TBI may demonstrate surprisingly few functional deficits. A goal of future research will be to further clarify the reasons for variability in outcomes among children with TBI.

References

- Achenbach, T. M. (1991). *Manual for the Child Behavior Checklist/4-18 and 1991 Profile*. Burlington, VT: University of Vermont, Department of Psychiatry.
- Anderson, P., Anderson, V., Northam, E., & Taylor, H. G. (2000). Standardization of the Contingency Naming Test: A measure of reactive flexibility. *Clinical Neuropsychological Assessment*, 1, 247-273.
- Anderson, V. A., Catroppa, C., Haritou, F., Morse, S., & Rosenfeld, J. V. (2005). Identifying factors contributing to child and family outcome 30 months after traumatic brain injury. *Journal of Neurology, Neurosurgery & Psychiatry*, 76, 401-408.
- Anderson, V. A., Morse, S. A., Klug, G., Catroppa, C., Haritou, F., Rosenfeld, J., & Pentland, L. (1997). Predicting recovery from head injury in young children: A prospective analysis. *Journal of the International Neuropsychological Society*, 3, 568-580.
- Barry, C. T., Taylor, H. G., Klein, S., & Yeates, K. O. (1996). The validity of neurobehavioral symptoms in children with traumatic brain injury. *Child Neuropsychology*, 2, 213-226.
- Beery, K. E. (1989). *Revised administration, scoring, and teaching manual for the Developmental Test of Visual-Motor Integration*. Cleveland, OH: Modern Curriculum Press.
- Bergman, L. R., & Trost, K. (2006). The person-oriented versus the variable-oriented approach: Are they complementary, opposites, or exploring different worlds? *Merrill-Palmer Quarterly*, 52, 601-632.
- Chadwick, O., Rutter, M., Brown, G., Shaffer, D., & Traum, M. (1981). A prospective study of children with head injuries: II. Cognitive sequelae. *Psychological Medicine*, 11, 49-61.
- Connell, A. M., Dishion, T. J., & Deater-Deckard, K. (2006). Variable- and person-centered approaches to the analysis of early adolescent substance use: Linking peer, family, and intervention effects with developmental trajectories. *Merrill-Palmer Quarterly*, 52, 421-448.
- Delis, D., Kramer, J., Kaplan, E., & Ober, B. (1986). *The California Verbal Learning Test: Children's version*. San Antonio, TX: Psychological Corporation.
- Dennis, M., Wilkinson, M., Koski, L., & Humphreys R. P. (1995). Attention deficits in the long term after childhood head injury. In S. H. Broman & M. E. Michel (Eds.), *Traumatic head injury in children* (pp. 165-187). New York: Oxford University Press.
- Ewing-Cobbs, L., Fletcher, J. M., Levin, H. S., Iovino, I., & Miner, M. E. (1998). Academic achievement and academic placement following traumatic brain injury in children and adolescents: A two-year longitudinal study. *Journal of Clinical and Experimental Neuropsychology*, 20, 769-781.
- Fay, G. C., Jaffe, K. M., Polissar, M. L., Liao, S., Rivara, J. B., & Martin, K. M. (1994). Outcome of pediatric traumatic brain injury at three years: A cohort study. *Archives of Physical Medicine and Rehabilitation*, 75, 733-741.
- Fletcher, J. M., Ewing-Cobbs, L., Miner, M. E., Levin, H. S., & Eisenberg, H. M. (1990). Behavioral changes after closed head injury in children. *Journal of Consulting and Clinical Psychology*, 58, 93-98.
- Francis, D. J., Fletcher, J. M., Stuebing, K. K., Lyon, G. R., Shaywitz, B. A., & Shaywitz, S. E. (2005). Psychometric approaches to the identification of LD: IQ and achievement scores are not sufficient. *Journal of Learning Disabilities*, 38, 98-108.
- Golden, C. J., Purish, A. D., & Hammeke, T. A. (1991). *Luria-Nebraska Neuropsychological Battery: Forms I and II*. Los Angeles: Western Psychological Services.

- Hannay, H. J., & Lezak, M. D. (2004). The neuropsychological examination: Interpretation. In M. D. Lezak, D. B. Howieson, & D. W. Loring (Eds.), *Neuropsychological assessment* (4th ed., pp. 133–156). New York: Oxford University Press.
- Heaton, R. K., Grant, I., & Matthews, C. G. (1991). *Comprehensive norms for an expanded Halstead-Reitan battery: Demographic corrections, research findings, and clinical applications*. Odessa, FL: Psychological Assessment Resources.
- Ingraham, L. J., & Aiken, C. B. (1996). An empirical approach to determining criteria for abnormality in test batteries with multiple measures. *Neuropsychology*, 10, 120–124.
- Jacobson, N. S., Roberts, L. J., Berns, S. B., & McGlinchey, J. B. (1999). Methods for defining and determining the clinical significance of treatment effects: Description, application, and alternatives. *Journal of Consulting and Clinical Psychology*, 67, 300–307.
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12–19.
- Jaffe, K. M., Polissar, N. L., Fay, G. C., & Liao, S. (1995). Recovery trends over three years following pediatric traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 76, 17–26.
- Kinsella, G., Ong, B., Murtaugh, D., Prior, M., & Sawyer, M. (1999). The role of the family for behavioral outcome in children and adolescents following traumatic brain injury. *Journal of Consulting and Clinical Psychology*, 67, 116–123.
- Kinsella, G., Prior, M., Sawyer, M., Murtagh, D., Eisenmajer, R., Anderson, V., et al. (1995). Neuropsychological deficit and academic performance in children and adolescents following traumatic brain injury. *Journal of Pediatric Psychology*, 20, 753–767.
- Knights, R. M., & Norwood, J. A. (1980). *Revised smoothed normative data on the Neuropsychological Test Battery for Children*. Ottawa, Ontario: Carleton University.
- Laursen, B., & Hoff, E. (2006). Person-centered and variable-centered approaches to longitudinal data. *Merrill-Palmer Quarterly*, 52, 377–389.
- Lindgren, S., & Lyons, D. (1984). *Pediatric Assessment of Cognitive Efficiency (PACE)*. Iowa City, IA: Department of Pediatrics, University of Iowa.
- Magnusson, D. (2003). The person approach: Concepts, measurement models, and research strategy. *New Directions for Child and Adolescent Development*, 101, 3–23.
- Massagli, T. L., Michaud, L. J., & Rivara, F. P. (1996). Association between injury indices and outcome after severe traumatic brain injury in children. *Archives of Physical Medicine and Rehabilitation*, 77, 125–132.
- Masten, A. S. (2001). Ordinary magic: Resilience processes in development. *American Psychologist*, 56, 227–238.
- Max, J. E., Robin, D. A., Lindgren, S. D., Smith, W. L., Jr., Sato, Y., Mattheis, P. J., et al. (1997). Traumatic brain injury in children and adolescents: Psychiatric disorders at two years. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36, 1278–1285.
- Mayer, T., Matlack, M., Johnson, D., & Walker, M. (1980). The Modified Injury Severity Scale in pediatric multiple trauma patients. *Journal of Pediatric Surgery*, 15, 719–726.
- McDonald, C. M., Jaffe, K. M., Fay, G. C., Polissar, N. L., Martin, K. M., Liao, S., & Rivara, J. B. (1994). Comparison of indices of traumatic brain injury severity as predictors of neurobehavioral outcome in children. *Archives of Physical Medicine and Rehabilitation*, 75, 328–337.
- Miller, I., Bishop, D., Epstein, N., & Keitner, G. (1985). The McMaster Family Assessment Device: Reliability and validity. *Journal of Marital and Family Therapy*, 11, 345–356.
- Moos, R., & Moos, B. (1994). *Life Stressors and Social Resources Inventory—Adult Form: Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Muthén, B., & Muthén, L. K. (2000). Integrating person-centered and variable-centered analyses: Growth mixture modeling with latent trajectory classes. *Alcoholism: Clinical and Experimental Research*, 24, 882–891.
- Perrot, S. B., Taylor, H. G., & Montes, J. L. (1991). Neuropsychological sequelae, family stress, and environmental adaptation following pediatric head injury. *Developmental Neuropsychology*, 7, 69–86.
- Rourke, B., & Orr, R. (1977). Prediction of the reading and spelling performances of normal and retarded readers: A four-year follow-up. *Journal of Abnormal Child Psychology*, 5, 9–20.
- Schwartz, L., Taylor, H. G., Drotar, D., Yeates, K. O., Wade, S. L., & Stancin, T. (2003). Long-term behavior problems following pediatric traumatic brain injury: Prevalence, predictors, and correlates. *Journal of Pediatric Psychology*, 28, 251–263.
- Semel, E., Wiig, E. H., Secord, W., & Sabers, D. (1987). *CELF-R: Clinical Evaluation of Language Fundamentals—Revised (technical manual)*. New York: Psychological Corporation.
- Slick, D. J. (2006). Psychometrics in neuropsychological assessment. In E. Strauss, E. M. S. Sherman, & O. Spreen (Eds.), *A compendium of neuropsychological tests: Administration, norms, and commentary* (3rd ed., pp. 3–43). New York: Oxford University Press.
- Sparrow, S., Balla, D., & Cicchetti, D. (1984). *Vineland Adaptive Behavior Scales*. Circle Pines, MN: American Guidance Service.
- Spreen, O., & Strauss, E. (1991). *A compendium of neuropsychological tests: Administration, norms, and commentary*. New York: Oxford University Press.
- Stancin, T., Drotar, D., Taylor, H. G., Yeates, K. O., Wade, S. L., & Minich, N. (2002). Health-related quality of life of children and adolescents after traumatic brain injury. *Pediatrics*, 109, 1–8.
- Stevens, G., & Featherman, D. L. (1981). A revised socioeconomic index of occupational status. *Social Science Research*, 10, 364–395.
- Stevenson-Hinde, J., & Akister, J. (1995). The McMaster Model of Family Functioning: Observer and parental ratings in a nonclinical sample. *Family Process*, 34, 337–347.
- Taylor, H. G. (2004). Research on outcomes of pediatric traumatic brain injury: Current advances and future directions. *Developmental Neuropsychology*, 25, 199–225.
- Taylor, H. G., Hack, M., & Klein, N. (1998). Attention deficits in children with <750 gm birth weight. *Child Neuropsychology*, 4, 21–34.
- Taylor, H. G., & Schatschneider, C. (1992). Child neuropsychological assessment: A test of basic assumptions. *The Clinical Neuropsychologist*, 6, 259–275.
- Taylor, H. G., Yeates, K. O., Wade, S. L., Drotar, D., Klein, S. K., & Stancin, T. (1999). Influences on first-year recovery from traumatic brain injury in children. *Neuropsychology*, 13, 76–89.
- Taylor, H. G., Yeates, K. O., Wade, S. L., Drotar, D., Stancin, T., & Burant, C. (2001). Bidirectional child–family influences on outcomes of traumatic brain injury in children. *Journal of the International Neuropsychological Society*, 7, 755–767.
- Taylor, H. G., Yeates, K. O., Wade, S. L., Drotar, D., Stancin, T., & Minich, N. (2002). A prospective study of short- and long-term outcomes after traumatic brain injury in children: Behavior and achievement. *Neuropsychology*, 16, 15–27.
- Taylor, H. G., Yeates, K. O., Wade, S. L., Drotar, D., Stancin, T., & Montpetite, M. (2003). Long-term educational interventions after traumatic brain injury in children. *Rehabilitation Psychology*, 48, 227–236.
- Teasdale, G., & Jennett, B. (1974). Assessment of coma and impaired consciousness: A practical scale. *Lancet*, 2, 81–84.
- von Eye, A., & Bogat, L. R. (2006). Person-oriented and variable-oriented research: Concepts, results, and development. *Merrill-Palmer Quarterly*, 52, 390–420.

- Wechsler, D. (1991). *WISC-III: Wechsler Intelligence Scale for Children—Third Edition*. San Antonio, TX: Psychological Corporation.
- Woodcock, R., & Mather, N. (1989). *Woodcock-Johnson Tests of Achievement—Revised—Standard and supplemental batteries*. Allen, TX: DLM Teaching Resources.
- Yeates, K. O. (1994). Comparison of developmental norms for the Boston Naming Test. *The Clinical Neuropsychologist*, 8, 91–98.
- Yeates, K. O., & Taylor, H. G. (1997). Predicting premorbid neuropsychological functioning following pediatric traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology*, 19, 825–837.
- Yeates, K. O., & Taylor, H. G. (2006). Behavior problems in school and their educational correlates among children with traumatic brain injury. *Exceptionality*, 14, 141–154.
- Yeates, K. O., Taylor, H. G., Barry, C., Drotar, D., Wade, S., & Stancin, T. (2001). Neurobehavioral symptoms in childhood closed-head injuries: Changes in prevalence and correlates during the first year postinjury. *Journal of Pediatric Psychology*, 26, 79–91.
- Yeates, K. O., Taylor, H. G., Wade, S. L., Drotar, D., Stancin, T., & Minich, N. (2002). A prospective study of short- and long-term neuropsychological outcomes after pediatric traumatic brain injury. *Neuropsychology*, 16, 514–523.

Received June 24, 2008

Revision received September 11, 2008

Accepted November 24, 2008 ■

Low Publication Prices for APA Members and Affiliates

Keeping you up-to-date. All APA Fellows, Members, Associates, and Student Affiliates receive—as part of their annual dues—subscriptions to the *American Psychologist* and *APA Monitor*. High School Teacher and International Affiliates receive subscriptions to the *APA Monitor*, and they may subscribe to the *American Psychologist* at a significantly reduced rate. In addition, all Members and Student Affiliates are eligible for savings of up to 60% (plus a journal credit) on all other APA journals, as well as significant discounts on subscriptions from cooperating societies and publishers (e.g., the American Association for Counseling and Development, Academic Press, and Human Sciences Press).

Essential resources. APA members and affiliates receive special rates for purchases of APA books, including the *Publication Manual of the American Psychological Association*, and on dozens of new topical books each year.

Other benefits of membership. Membership in APA also provides eligibility for competitive insurance plans, continuing education programs, reduced APA convention fees, and specialty divisions.

More information. Write to American Psychological Association, Membership Services, 750 First Street, NE, Washington, DC 20002-4242.