

Terminal Decline in Well-Being: The Role of Social Orientation

Denis Gerstorf

Humboldt University and German Institute for Economic Research (DIW Berlin), Berlin, Germany

Christiane A. Hoppmann

University of British Columbia

Corinna E. Löckenhoff

Cornell University

Frank J. Infurna

Arizona State University

Jürgen Schupp

German Institute for Economic Research (DIW Berlin), Berlin, Germany, and Free University, Berlin

Gert G. Wagner

Max Planck Institute for Human Development, Berlin, and German Institute for Economic Research (DIW Berlin), Berlin, Germany

Nilam Ram

Pennsylvania State University and German Institute for Economic Research (DIW Berlin), Berlin, Germany

Well-being development at the end of life is often characterized by steep deteriorations, but individual differences in these terminal declines are substantial and not yet well understood. This study moved beyond typical consideration of health predictors and explored the role of social orientation and engagement. To do so, we used social variables at the behavioral level (self-ratings of social participation) and the motivational level (valuing social and family goals), assessed 2 to 4 years before death. We applied single- and multiphase growth models to up to 27-year annual longitudinal data from 2,910 now deceased participants of the nation-wide German Socio-Economic Panel Study ($M_{\text{age at death}} = 74$ years; $SD = 14$; 48% women). Results revealed that leading a socially active life and prioritizing social goals in late life were independently associated with higher late-life well-being, less pronounced late-life decline, and a later onset of terminal decline. Significant interaction effects suggested that the combination of (reduced) social participation and (lowered) social goals magnifies the effects of each other. Findings also indicated that less decline in social participation was associated with less severe rates and a later onset of well-being decline. We found little evidence that valuing family goals is associated with late-life trajectories of well-being. Associations were independent of key correlates of well-being and mortality, including age at death, gender, education, disability, hospital stays, and goals in other life domains. We discuss possible pathways by which maintaining social orientation into late life may help mitigate terminal decline in well-being.

Keywords: successful aging, life satisfaction, social support, development, German Socio-Economic Panel Study

A central objective of developmental research is to describe and explain interindividual differences in intraindividual change (Baltes & Nesselrode, 1979). One of the most intriguing phenomena

in this regard are changes in central areas of life, including well-being, that occur at the end of life (i.e., terminal decline; Burns, Mitchell, Shaw, & Anstey, 2014; Holt-Lunstad, Smith, & Layton,

Editor's Note. Elliot M. Tucker-Drob served as the action editor for this article.—UM

Denis Gerstorf, Institute for Psychology, Humboldt University, and German Institute for Economic Research (DIW), Berlin, Germany; Christiane A. Hoppmann, Department of Psychology and Center for Hip Health and Mobility, University of British Columbia; Corinna E. Löckenhoff, Department of Human Development, Cornell University; Frank J. Infurna, Department of Psychology, Arizona State University; Jürgen Schupp, German Institute for Economic Research (DIW), Berlin, Germany, and Department of Political and Social Sciences, Free University, Berlin; Gert G. Wagner, Max Planck Institute for Human Development, Berlin, and German Institute for Economic Research (DIW Berlin), Berlin, Germany; and Nilam Ram, Department of

Human Development and Family Studies, Pennsylvania State University, and German Institute for Economic Research (DIW), Berlin, Germany.

Supported by the German Research Foundation (Grant GE 1896/3-1); the National Institute on Aging (Grants RC1-AG035645, R01-AG048844, R21-AG032379, and R21-AG033109); the Max Planck Society and the Max Planck Institute for Human Development, Berlin; the Penn State Children, Youth, and Families Consortium; Christiane Hoppmann acknowledges the support of the Michael Smith Foundation for Health Research and the Canada Research Chairs program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

Correspondence concerning this article should be addressed to Denis Gerstorf, Humboldt University Berlin, Institute of Psychology, Rudower Chaussee 18, 12489 Berlin, Germany. E-mail: denis.gerstorf@hu-berlin.de

2010; Mroczek & Spiro, 2005; Schilling, Wahl, & Wiegering, 2013). Although individual differences in terminal decline in well-being are well documented and substantial, the specific factors contributing to such differences are not yet fully understood, particularly outside the physical health domain (see Gerstorf & Ram, 2013, for overview). The current study drew on a tradition of research implicating social orientation and social resources as fundamental components of successful aging (Rowe & Kahn, 1997) and examined whether and how social orientation is associated with differences in terminal decline in well-being. To do so, we used social variables at the behavioral level (self-ratings of social participation) and the motivational level (valuing social and family goals), obtained an average of 2–4 years before death. We applied single- and multiphase growth models to up to 27-year annual longitudinal data from 2,910 now-deceased participants of the nation-wide German Socio-Economic Panel Study (SOEP) and examined whether and how social orientation is associated with late-life well-being levels, rates of terminal decline, and delayed onset of terminal decline. To control for known correlates of well-being and mortality, all our models covaried for age at death, gender, education, disability, hospital stays, and goals in other life domains.

Well-Being Trajectories in Adulthood and Late Life

Theories of self-regulation and myriad empirical reports attest to the relative stability of well-being across adulthood and into old age (Carstensen et al., 2011; Charles, Reynolds, & Gatz, 2001; Diener, Lucas, & Scollon, 2006; Ryff & Singer, 1998). Qualifying this overall positive picture, a growing body of research has repeatedly shown that mortality-related time-to-death models typically account for more variance in interindividual differences in late-life well-being change than age-related models and that, on average, both cognitive evaluative and emotional well-being decline rapidly in the last years of life (see Gerstorf & Ram, 2013, for overview). To use an example from our own work, participants across three national studies in the United States, Great Britain, and Germany were found to enter the terminal phase of steep well-being deteriorations an average of 3–5 years prior to death (Gerstorf et al., 2010). To illustrate, the average decedent from the German SOEP study exhibited almost a full standard deviation (effect size units) decline in life satisfaction during the last 4 years of life. However, despite this overall negative picture of terminal decline in well-being, considerable individual differences exist (Burns, Byles, Magliano, Mitchell, & Anstey, 2015; Gerstorf & Ram, 2013). Although many people exhibit dramatic declines in well-being in the last years of their lives, some individuals are able to maintain their well-being until (or very close to) the end of life.

Previous evidence has shown, through proxy variables, that pathology, disability, and comorbidities contribute to late-life well-being (Burns et al., 2014; Gerstorf, Ram, Lindenberger, & Smith, 2013; Infurna et al., 2014). Interestingly, these proxies are often associated with differences in level of well-being late in life, but not associated with differences in rate of terminal decline. Setting limitations in statistical power aside, these findings suggest that *biological factors* (e.g., poor health) are one of several sets of factors that shape late-life well-being. In line with the notion that terminal decline is not just biological, recent evidence has demonstrated that *psychological resources* (e.g., perceived personal control) are also associated with higher levels of late-life well-

being, less severe rates of terminal decline, and later onset of terminal decline (Gerstorf et al., 2014). Continuing in the quest to help vulnerable segments of the population by identifying modifiable psychosocial factors related to differences in terminal decline in well-being, the current study examined the role of *psychosocial functioning*, namely, social orientation.

Social Orientation and Well-Being

Social relationships and activities have long been identified as key elements of successful aging (Rowe & Kahn, 1997, 2015) and there is solid evidence that they contribute to individual health and well-being (Berkman, Glass, Brissette, & Seeman, 2000; Cohen, 2004; Diener & Seligman, 2004; Uchino, 2006). For example, Antonucci (2001) concluded from her literature overview that people who report being socially integrated also tend to report higher well-being. Conversely, social isolation was found to be associated with lower mental and physical health, especially among older adults (Dahlberg & McKee, 2014; Hawton et al., 2011; Ong, Rothstein, & Uchino, 2012). In the current study, we focused on process aspects of the Fingerman and Lang's (2004) framework of social relations and specifically examined two aspects of social orientation and engagement: (a) at the behavioral level, social activities and participation, and (b) at the motivational level, the importance of social and family goals.

Social Activities and Participation

Going back to influential notions of activity theory, according to which social involvement and activity constitute prerequisites for successful aging (Burgess, 1954; Lemon, Bengtson, & Peterson, 1972), social engagement is considered a key driver of life satisfaction in older adults. Social activity may influence well-being directly, for example, through the pursuit of kind acts and engagement in meaningful and joyful activities (Lyubomirsky & Layous, 2013). The association may also be more indirect, for example, in that social activities promote feelings of competence, physical health, and cognitive functioning (Herzog & House, 1991; Menec, 2003; Netz, Wu, Becker, & Tenenbaum, 2005), which in turn contribute to high well-being. Empirical evidence has largely supported these propositions, both in cross-sectional data (Herzog, Franks, Markus, & Holmberg, 1998; Lawton, Ruckdeschel, Winter, & Kleban, 1999; Litwin & Shiovitz-Ezra, 2006) and longitudinally (Menec, 2003; Sneed & Cohen, 2013). For example, Huxhold, Fiori, and Windsor (2013) recently reported that social engagement among older adults was predictive of increases in or maintenance of life satisfaction across 6 years.

Social and Family Goals

At the level of goals, we followed Headey (2008) and distinguished broad social goals geared toward diverse social relationships and general social participation from those geared toward core family members (e.g., spouses, children). Beginning with the former, *general social and altruistic goals* (e.g., engaging in volunteering activities; Headey, Muffels, & Wagner, 2010) are important for well-being for a several reasons. For one, social goals concerning one's ability to help other people can facilitate self-esteem, a sense of control, and feelings of competence, for example,

through confirming role identities and self-concepts (Glass, Seeman, Herzog, Kahn, & Berkman, 1995; Herzog & House, 1991), and fostering a sense of generativity (Erikson, 1985). In a similar vein, maintaining involvement in social and political activities, despite the challenges that often accompany old age, may convey the self-protective impression of a certain degree of continuity with earlier phases of life (Atchley, 1989). Like broader social goals, *family-related goals* may contribute to older adults' well-being by maintaining roles, identities, and self-esteem (Mutran, 1987). Moreover, the active pursuit of family goals promotes access to a network of close confidants, which was found to be a strong predictor of well-being in older populations (Litwin & Stoeckel, 2014). According to socioemotional selectivity theory (SST; Carstensen, 1995, 2006; Lang & Carstensen, 2002), age-related limitations in perceived time left in life trigger a motivational shift toward a preference for close social partners (Fredrickson & Carstensen, 1990; Fung, Carstensen, & Lutz, 1999; Lang & Carstensen, 1994), suggesting that family-related goals are particularly important for older individuals. Taken together, the previous literature has provided considerable evidence for the benefits of social goals and social participation across the life span and into old age. However, this work has largely drawn on fairly healthy, community-dwelling samples and it is not entirely clear whether such effects are maintained as individuals enter the phase of terminal decline.

Social Orientation and Well-Being at the End of Life

Although the basic pattern of positive associations between social engagement and well-being is likely to continue into the final years of life, some theoretical perspectives would suggest that effects may become attenuated or more complex. With regard to social participation, for example, decades of research on cancer support groups have been built on the notion that social engagement is a key resource when facing potentially life-threatening health conditions (Rehse & Pukrop, 2003). However, some have argued that the benefits of social support are limited when stress levels are high (Taylor, 2011), as is often the case at the end of life. Moreover, maintaining high levels of social participation at all cost could represent a strain on dying individuals' increasingly limited resources and thus present a detriment to well-being (Charles, 2010). Similarly, with regard to goal priorities, the pursuit of broad and altruistic social goals at the end of life may continue to convey benefits for self-esteem and feelings of generativity, but it could also interfere with the reorientation toward close social relationships in the face of limited time that has been proposed by SST (Lang, 2000).

Conceivably, end-of-life trajectories in social goals and social participation may also interact with each other. According to the motivational theory of life span development (Heckhausen, Wrosch, & Schulz, 2010), active goal pursuit is only beneficial for well-being if the goal in question is still within reach. Thus, individuals who retain high levels of general social goals but lack the resources for active social participation could fare worse than those who adaptively disengage from such goals (Heckhausen, Wrosch, & Schulz, 2013). On the other hand, there is some evidence that maintaining a sense of engagement with family and friends can benefit dying individuals' mental well-being, even if active engagement in such relationships is increasingly curtailed (Prince-Paul, 2008). Thus, it is not clear whether active social participation and the pursuit of social goals affect well-being

independently, magnify the effects of one another, or interact in more complex ways.

The Present Study

In the current report, we examined whether and how aspects of social orientation (i.e., self-reports of engaging in social activities and prioritizing social and family goals) shape terminal decline in well-being. To this end, we applied single- and multiphase models of change to long-term longitudinal data from now-deceased participants in Germany's nation-wide SOEP study. We expected that both higher levels and more favorable trajectories in social orientation would be associated with higher late-life well-being as well as later onset and less severe rates of late life decline. We also explored possible interactions between social goals and social activities to gather insights into the complex nature of associations between these variables. Our models covaried for age at death, gender, education, disability, hospital stays, and goals in other life domains. These variables are important to take into account because, for example, older age, being a man, little education, and poor health have each been associated with poorer well-being and increased mortality hazards (Antonucci, 2001; Burns et al., 2015; DeNeve & Cooper, 1998; Guralnik, 1991; Hoppmann, Gerstorf, Smith, & Klumb, 2007; Moen, 2001).

Method

To examine our research questions, we used data from the German SOEP study, which is increasingly used in psychological research (see Siedler, Schupp, & Wagner, 2011, for overview). Information about the design, participants, variables, and assessment procedures in the larger study has been reported in Headey et al. (2010). A brief overview of details relevant to the present analysis is given below.

Participants and Procedure

The SOEP is a nationally representative annual longitudinal panel study of private households covering currently about 50,000 adult residents of former West and East Germany, including immigrants and resident foreigners. Potential participants were drawn at random from a set of randomly selected geographic locations in Germany. Initial response rates of 60%–70% and relatively low longitudinal attrition (about 15% for the second wave and less than 5% yearly attrition later) resulted in an overall sample that was shown to be representative of the population living in private households (Kroh, Pischner, Spieß, & Wagner, 2008) and long-term care homes in Germany (Klein, 1996). Data were predominantly collected in face-to-face interviews, and about 10% of long-term participants self-administered the questionnaires.

To examine terminal decline, we used information about mortality status and year of death for deceased participants that was obtained at the yearly interviews, either directly from the remaining household members or neighbors or from official registries. Given that death rates and ages of death of SOEP participants parallel official life tables, the SOEP has long served as a representative resource for mortality-related analyses in Germany (e.g., Brockmann & Klein, 2004; Burkhauser, Giles, Lillard, & Schwarze, 2005). Here, we analyzed data from the 2,910 partici-

pants who had (a) died prior to May 2011 and (b) had provided data on well-being, all social resource variables, and all covariates. Respondents included in our sample were born between 1894 and 1990 and died between 1991 and 2011. Acknowledging reports that terminal declines in well-being are not specific to old age (Gerstorff et al., 2010), we considered all by now deceased SOEP participants including 14% who died before age 60. Participants included in our analyses contributed an average of 11.75 ($SD = 6.19$) yearly reports, for a total of 33,777 observations. The large majority of participants provided one or more well-being ratings in the last 3 years of life (86%) and contributed five or more annual well-being observations in total (91%).

The primary reason for study dropout among older participants in the larger SOEP study was mortality, as opposed to other reasons (e.g., poor health, relocation, loss of interest; Kroh, 2014). To characterize our current analysis sample, we conducted three sets of selectivity analyses. First, as expected, compared with still-living participants ($n \approx 46,000$), the deceased subsample we used was, on average, older at their very first assessment in the SOEP ($M = 61.29$ years, $SD = 14.18$, vs. $M = 39.30$ years, $SD = 17.48$), $F(1, 49,011) = 4,268.3$, $R^2 = .080$; included more men (52% vs. 49%), $\chi^2 = (1, N = 49,015) = 13.1$; had fewer years of education ($M = 10.88$ years, $SD = 2.11$, vs. $M = 11.42$ years, $SD = 2.65$), $F(1, 49,013) = 846.2$, $R^2 = .003$; and reported lower levels of life satisfaction at their first wave of assessment ($M = 7.04$, $SD = 2.20$, vs. $M = 7.32$, $SD = 1.90$) on a 0–10 scale, $F(1, 49,013) = 59.2$, $R^2 = .001$; $ps < .001$. Second, we examined whether and how deceased SOEP participants included in our analyses differed from other deceased SOEP participants who had not been included (e.g., because of missing data on the relevant social orientation variables; $n \approx 2,100$). These analyses revealed that participants included in our subsample were younger at their first SOEP assessment (61.29 years, $SD = 14.18$, vs. 65.64 years, $SD = 15.49$), $F(1, 5,050) = 107.0$, $R^2 = .021$; had more years of education ($M = 10.88$ years, $SD = 2.11$, vs. $M = 10.49$ years, $SD = 2.10$), $F(1, 4,966) = 40.9$, $R^2 = .008$; and reported higher levels of life satisfaction at their first wave of assessment ($M = 7.04$, $SD = 2.20$, vs. $M = 6.80$, $SD = 2.43$) on a 0–10 scale, $F(1, 5,050) = 69.21$, $R^2 = .003$, $ps < .001$, whereas no differences were found for gender. Finally, examining sample attrition over time within our subsample of deceased SOEP participants, we compared those providing more data points (five and more waves: $n = 2,635$) with those providing fewer data points (four and fewer waves: $n = 275$). Analyses revealed that providing more data points was associated with younger age at first wave of assessment ($M = 60.87$ years, $SD = 13.83$, vs. $M = 65.34$ years, $SD = 16.62$), $F(1, 2,909) = 24.9$, $R^2 = .009$; and with higher levels of life satisfaction at the first wave of assessment ($M = 7.12$, $SD = 2.18$, vs. $M = 6.25$, $SD = 2.27$) on a 0–10 scale, $F(1, 2,909) = 39.3$, $R^2 = .013$, $ps < .001$, whereas no differences were found for gender and education.

Measures

Well-being. Life satisfaction was measured as responses to the question: “How satisfied are you with your life concurrently, all things considered?” (in German: “Wie zufrieden sind Sie gegenwärtig, alles in allem, mit ihrem Leben?”), answered on a scale from 0 (*totally unsatisfied*) to 10 (*totally satisfied*). This item, asked at yearly intervals, is considered a measure of cognitive–evaluative (as opposed to affective) well-being and has been widely used in psychological

research (see Fujita & Diener, 2005; Lucas, Clark, Georgellis, & Diener, 2003; Headey et al., 2010, for measurement properties of the item). On average, deaths occurred 2.07 years ($SD = 2.66$) after the last well-being rating. For consistency with other reports, life satisfaction scores were standardized to a T metric ($M = 50$, $SD = 10$) using the 2002 SOEP sample as the reference frame ($M = 6.90$, $SD = 1.81$; scale range: 0–10).

Social orientation. To index social orientation, we used measures that tap into social activities and participation and that tap into the importance of general social goals and more narrow family goals. With our interest in better understanding the actual (social) resources available and the constraints people face late in life, we indexed the degree of social orientation by using the last available observation (i.e., closest to death). By contrast, if our primary interest had been in the resources and constraints people bring into late life, we could have used the first wave of assessment for the social orientation variables. Thus, results obtained in this report may not necessarily generalize to social orientation variables as tracked with other operational definitions and within other lifetime periods. Probable late-life declines in social resources would make our approach a more conservative test of their protective role against late-life decrements in well-being.

Participant’s *social activities and participation* were measured using items asking the frequency (1 = *at least once a week*, 2 = *at least once a month*, 3 = *seldom*, and 4 = *never*) of involvement in or attendance at social networking and community activities: (a) visit cultural events such as concerts, theaters, or lectures, (b) active sport participation, (c) honorary activities in clubs, organizations, or social service, and (d) participation in citizen initiatives, parties, community politics. The scale is highly similar in structure to other measures of social participation (see Parslow, Jorm, Christensen, & Mackinnon, 2006). The social participation items endorsed most frequently were sport activities and honorary/cultural activities. Items were reverse coded and averaged to obtain an index with higher scores indicating more social participation. Social participation was assessed in 1985, 1986, 1988, 1990, 1992, 1994, 1995, 1996, 1997, 1998, 1999, 2001, 2003, 2005, 2007, 2008, 2009, and 2011. As above, we used of the last available social participation rating (i.e., closest to death). These ratings were obtained, on average, 2.46 years ($SD = 2.74$) before death. Details about the measurement properties of the social activities measure as used in the SOEP can be obtained from Infurna, Gerstorff, Ram, Schupp, and Wagner (2011).

The importance of general *social goals* was measured using items asking about the importance of two altruistic non-zero sum goals: “Various things can be important for various people. Are the following things currently . . . very important/important/less important or/not at all important . . . for you?”: (a) helping other people and (b) being involved in social and political activities. In a similar vein, the importance of *family goals* was assessed by responses to two items asking how much importance (1 = *not at all important* and 4 = *very important*) the respondent attached to the goals of (a) having a good marriage and (b) having a good relationship with children. Life goals/priorities were measured five times between 1990 and 2008 (in 1990, 1992, 1995, 2004, and 2008). The last available goal importance rating was obtained an average of 4.08 years ($SD = 3.35$) before death. Details about measurement properties and factor structure of the goal measures as used in the SOEP can be found in Headey (2008).

Covariates. Various covariates were used to control for individual differences in nonsocial factors known to be related to well-being and mortality (i.e., age at death, gender, education, disability, hospital stays, and goals in other life domains). Using individuals' most recent reports, *years of completed education* were noted as the minimum number of years an individual needed to acquire his or her particular degree. In addition, our models included two key health factors as covariates. First, *disability* was assessed at each wave with a single item asking participants whether they had been "officially certified as having a reduced capacity to work or being severely handicapped" (see Lucas, 2007, for details). Thus, disability indicators were based on self-reports, but referred to official certifications. Our measure contrasts all participants who had been disabled at some point during the study ($n = 1,387$) with those who were not ($n = 1,523$). Second, *hospital stay last year* was indexed by responses to the question; "Were you ever admitted to a hospital for at least one night in year X (e.g., 2002)?" Data for the number of hospital stays last year were taken from the last available data point from each participant. Assessments of hospital stays last year were obtained an average of 2.15 years ($SD = 2.71$) before death. Finally, to contrast the importance of social and family goals with those in other domains of life, we also included goal commitment ratings of other domains into our analyses. Specifically, we considered material success goals (being able to buy things, being successful in one's career), each rated using a 5-point Likert scale, ranging from 1 (*not at all important*) to 4 (*very important*). Again, we used the last available goal importance rating from each participant.

Intercorrelations for the variables we studied are reported in Table 1. We found that social participation, social goals, and family goals correlated between .07 and .33, indicating that these tap into distinct aspects of social orientation. Table 1 also indicates that the covariates interrelated with well-being at the last observation before death (e.g., $r = -.20$ for hospital stays) and with the social orientation measures (e.g., $r = .32$ between social participation and education), signifying the importance of the covariates.

Data Preparation

Using a multilevel modeling framework, growth curve models (e.g., McArdle & Nesselroade, 2003; Ram & Grimm, 2015; Singer & Willett, 2003) were used to summarize and extract information about rates of change in social participation, social goals, and family goals. Models took the following form;

$$\text{Social orientation}_{it} = \beta_{0i} + \beta_{1i}(\text{time}_{it}) + e_{it}, \quad (1)$$

where person i 's score in a particular social orientation variable at time t , $\text{social orientation}_{it}$, is a function of an individual-specific intercept parameter, β_{0i} , and an individual-specific linear slope parameter, β_{1i} , that captures the linear rate of change per year of time, and residual error, e_{it} . Following standard growth curve modeling procedures, individual-specific intercepts, β_{0i} , and linear slopes, β_{1i} (from the Level 1 model give in Equation 1), were modeled as:

$$\beta_{0i} = \gamma_{00} + u_{0i}, \quad (2)$$

$$\beta_{1i} = \gamma_{10} + u_{1i},$$

(i.e., Level 2 model) where γ_{00} and γ_{10} are sample means, and u_{0i} and u_{1i} are individual deviations from those means. Using SAS Proc Mixed with restricted maximum likelihood estimation and standard missing at random assumptions (Little & Rubin, 1987), we fitted the model separately for each of the three social orientation measures. Using Bayes empirical estimates (see Littell, Milliken, Stroup, Wolfinger, & Schabenberger, 2006), we obtained *rate of change*, β_{1i} , scores for each domain for each individual. Our objective was to reduce the longitudinal, up to 27 years, data across the three social orientation dimensions (social participation, social goals, and family goals) down to three informative scores about longitudinal change. As noted in the Measures section, we used the last available (i.e., closest to death) observation before death to represent the level of social orientation.

Data Analysis

To examine our research questions, we estimated both single- and multiphase growth models. In a first set of models, we fitted

Table 1
Intercorrelations Among Study Variables

Variables	<i>M</i>	<i>SD</i>	Intercorrelations										
			1	2	3	4	5	6	7	8	9	10	
Well-being													
1. Life satisfaction (<i>T</i> score)	43.29	13.66		.22	.17	.03	.05	-.02	.08	-.20	-.20	.07	
Social orientation													
2. Social participation (1-4)	1.28	0.45			.33	.07	-.25	-.10	.32	-.10	-.10	.19	
3. Social goals (1-4)	2.33	0.55				.23	-.13	-.10	.23	-.02	-.03	.25	
4. Family goals (1-4)	3.15	0.83					-.13	-.15	.06	.04	.02	.30	
Covariates													
5. Age at death (20-102 years)	74.12	13.66						.19	-.11	.02	-.01	-.35	
6. Women (0-1)	0.48	0.50							-.22	-.10	-.05	-.19	
7. Education (7-18)	10.88	2.11								-.01	-.04	.14	
8. Disabled (0-1)	0.48	0.50									.25	-.03	
9. Hospital stay last year (0-1)	0.19	0.39											-.00
10. Material success goals (1-4)	2.53	0.72											

Note. $N = 2,910$ participants. Life satisfaction, last observation before death used. Intercorrelations of $r = |.06|$ or above differ statistically significantly from zero at p greater than .001. Scores for life satisfaction were standardized to a *T* metric ($M = 50$; $SD = 10$) using the 2002 German Socio-Economic Panel Study sample as the reference frame ($M = 6.90$, $SD = 1.81$; scale range: 0-10).

single-phase growth curve models for well-being over time to death to effectively model between-person differences in how individuals' well-being changed with impending mortality. This model was specified as:

$$\text{Well-being}_{it} = \beta_{0i} + \beta_{2i}(\text{time to death}_{it}) + \beta_{4i}(\text{time to death}_{it}^2) + e_{it}, \quad (3)$$

where person *is* well-being at occasion *t*, well-being_{*it*}, is a function of an individual-specific intercept parameter, β_{0i} , individual-specific linear and quadratic slope parameters, β_{2i} and β_{4i} , and residual error, e_{it} . Following standard multilevel and latent growth modeling procedures (Ram & Grimm, 2015; Singer & Willett, 2003), individual-specific intercepts, β_{0i} , and slopes, β_{2i} and β_{4i} (from the Level 1 model given in Equation 3) were modeled as a function of the social orientation variables and the covariates. To facilitate model parsimony and convergence, nonsignificant interaction terms were trimmed, always retaining the lower order interactions where necessary (e.g., see Equation 6 below). Note that the starting models included both level of social orientation and change in social orientation as predictors (obtained as described above) in each (Level 2) equation, and were then trimmed to retain only those effects of change in social orientation that were statistically significant (i.e., change in social participation noted in Equations 4–6; level was retained consistently). The final model took the following form:

$$\begin{aligned} \beta_{0i} = & \gamma_{00} + \gamma_{01}(\text{age at death}_i) + \gamma_{02}(\text{women}_i) + \gamma_{03}(\text{education}_i) \\ & + \gamma_{04}(\text{disability}_i) + \gamma_{05}(\text{hospital}_i) \\ & + \gamma_{06}(\text{material success goals}_i) + \gamma_{07}(\text{social participation}_i) \\ & + \gamma_{08}(\text{social goals}_i) + \gamma_{09}(\text{family goals}_i) \\ & + \gamma_{10}(\text{social participation}_i * \text{social goals}_i) \\ & + \gamma_{11}(\text{change social participation}_i) + u_{0i}, \end{aligned} \quad (4)$$

$$\begin{aligned} \beta_{2i} = & \gamma_{20} + \gamma_{21}(\text{age at death}_i) + \gamma_{22}(\text{women}_i) + \gamma_{23}(\text{education}_i) \\ & + \gamma_{24}(\text{disability}_i) + \gamma_{25}(\text{hospital}_i) \\ & + \gamma_{26}(\text{material success goals}_i) + \gamma_{27}(\text{social participation}_i) \\ & + \gamma_{28}(\text{social goals}_i) + \gamma_{29}(\text{family goals}_i) \\ & + \gamma_{30}(\text{social participation}_i * \text{social goals}_i) \\ & + \gamma_{31}(\text{change social participation}_i) + u_{2i}, \end{aligned} \quad (5)$$

$$\beta_{4i} = \gamma_{40} + \gamma_{41}(\text{hospital}_i) + \gamma_{42}(\text{social participation}_i) + u_{4i}, \quad (6)$$

where the γ s are sample-level associations, and u_{0i} , u_{2i} , and u_{4i} are residual unexplained individual differences that are assumed to be multivariate normally distributed, correlated with each other, and uncorrelated with the residual errors, e_{it} . Cubic terms were also included and tested, but did not differ significantly from zero and were thus not included in the final models. The time-to-death variable was centered at 1 year prior to death. All predictors were effect-coded/centered so that the regression parameters for these variables indicated the average trajectory and the extent of differences associated with a particular variable (rather than for a particular group). Negative parameters indicated that a given social resource (e.g., participation) or predictor variable (e.g., disability)

was associated with lower levels, steeper decline, or steeper curvature.

In a second set of models, we invoked notions of terminal decline by using extensions (Cudeck & Harring, 2007; Cudeck & Klebe, 2002) of multiphase or “spline” growth models (Ram & Grimm, 2015; Singer & Willett, 2003). To facilitate model parsimony and convergence, nonsignificant interaction terms, predictors, and some random effects were trimmed, always retaining the lower order interactions where necessary, and prioritizing those random effects of greatest substantive interest (e.g., terminal phase). For example, we removed the material success goals variable (which carried no predictive value in earlier models)¹, and all predictors for rates of preterminal decline [Equation 10]), and random effect for the transition point (i.e., no u_{6i} in Equation 12). As well, convergence was facilitated by parameterizing the random effect variance–covariance matrix using (squared) standard deviations. Again, although the change in social orientation variables was trimmed, the level of social orientation variables was consistently retained. The final model took the following form:

$$\begin{aligned} \text{Well-being}_{it} = & \beta_{0i} + \beta_{2i}(\text{time to death}_{it} - k_i) \\ & + e_{it}, \text{ when time to death}_{it} < k_i, \text{ and} \end{aligned} \quad (7)$$

$$\begin{aligned} \text{Well-being}_{it} = & \beta_{0i} + \beta_{4i}(\text{time to death}_{it} - k_i) \\ & + e_{it}, \text{ when time to death}_{it} \geq k_i, \end{aligned} \quad (8)$$

where β_{0i} represents a person *is* level of life satisfaction at k_i , a person-specific transition point that indicates a shift from the preterminal phase to the terminal phase; rate of change in the preterminal phase is captured by β_{2i} , and individual-specific rates of change in the terminal-phase by β_{4i} (time to death_{*it*} is coded in negative integers). Individual differences in the intercept β_{0i} (located at the time of the transition), slopes β_{2i} and β_{4i} , and the transition point k_i , were modeled as a function of the social orientation variables and covariates,

$$\begin{aligned} \beta_{0i} = & \gamma_{00} + \gamma_{01}(\text{age at death}_i) + \gamma_{02}(\text{women}_i) + \gamma_{03}(\text{education}_i) \\ & + \gamma_{04}(\text{disability}_i) + \gamma_{05}(\text{hospital}_i) + \gamma_{06}(\text{social participation}_i) \\ & + \gamma_{07}(\text{social goals}_i) + \gamma_{08}(\text{family goals}_i) \\ & + \gamma_{09}(\text{social participation}_i * \text{social goals}_i) \\ & + \gamma_{10}(\text{change social participation}_i) + u_{0i}, \end{aligned} \quad (9)$$

$$\beta_{2i} = \gamma_{20} + u_{2i}, \quad (10)$$

$$\begin{aligned} \beta_{4i} = & \gamma_{40} + \gamma_{41}(\text{age at death}_i) + \gamma_{42}(\text{women}_i) + \gamma_{43}(\text{education}_i) \\ & + \gamma_{44}(\text{disability}_i) + \gamma_{45}(\text{hospital}_i) + \gamma_{46}(\text{social participation}_i) \\ & + \gamma_{47}(\text{social goals}_i) + \gamma_{48}(\text{family goals}_i) \\ & + \gamma_{49}(\text{social participation}_i * \text{social goals}_i) \\ & + \gamma_{50}(\text{change social participation}_i) + u_{4i}, \end{aligned} \quad (11)$$

$$\begin{aligned} k_i = & \gamma_{60} + \gamma_{61}(\text{age at death}_i) + \gamma_{62}(\text{women}_i) + \gamma_{63}(\text{education}_i) \\ & + \gamma_{64}(\text{disability}_i) + \gamma_{65}(\text{hospital}_i) \end{aligned}$$

¹ In follow-up analyses, it was possible to estimate a model that had the family goals variable replaced with the material goals variable, everything else being equal. These analyses corroborated the main findings according to which social participation and social goals were independently associated with both higher late-life well-being, less pronounced late-life decline, and a later onset of terminal decline.

$$\begin{aligned}
& + \gamma_{66}(\text{social participation}_i) + \gamma_{67}(\text{social goals}_i) \\
& + \gamma_{68}(\text{family goals}_i) \\
& + \gamma_{69}(\text{social participation}_i * \text{social goals}_i) \\
& + \gamma_{70}(\text{change social participation}_i),
\end{aligned}$$

where the γ s are sample-level associations, and u_{0i} , u_{2i} , and u_{4i} are residual unexplained individual differences that are assumed to be multivariate normally distributed, correlated with each other, and uncorrelated with the residual errors, e_{it} . Paralleling interpretation of other parameters, negative parameters in Equation 12 indicate that a given social resource (e.g., participation) or predictor variable (e.g., disability) was associated with spending fewer years in the terminal decline phase (i.e., being closer to zero).

All models were fit to the data using SAS Proc Mixed (single-phase models) or Proc NLMixed (multiphase models; Littell et al., 2006), with incomplete data accommodated under usual missing at random assumptions (Little & Rubin, 1987). The predictors and covariates (e.g., age, health) represent attrition-informative variables and so helped to accommodate longitudinal selectivity for the outcome variable of well-being (McArdle, 1994). With our relatively large sample size, significance tests were evaluated at p less than .001.

Results

Does Social Orientation Relate to Higher Late-Life Well-Being and Less Severe Rates of Decline?

In a first set of analyses, we used single-phase models of change to examine whether social orientation and the covariates relate to higher late-life well-being (1 year prior to death) and less severe rates of decline. Results are reported in Table 2. Consistent with earlier work, we found that the typical trajectory of late-life well-being in this sample was characterized by linear decline with some acceleration. Specifically, the linear component of decline amounted to almost a full standard deviation per 10 years ($\gamma_{20} = -0.90$), which together with some concave curvature ($\gamma_{40} = -0.03$) brought the average individual to a well-being level at 1 year prior to death ($\gamma_{00} = 45.45$) that was 0.5 SD below the mean of the nationally representative SOEP sample in 2002 ($M = 50$, $SD = 10$). As expected, several of the covariates were associated with levels and rates of decline in well-being. Individuals who had died at older ages reported higher well-being in the year prior to death ($\gamma_{01} = 0.09$), probably a reflection of selection processes. Women reported lower well-being close to death ($\gamma_{02} = -1.32$). Individuals with disability had lower levels of well-being close to death ($\gamma_{04} = -3.46$) and experienced more precipitous declines ($\gamma_{24} = -0.10$). Similarly, those who had to spend one night or more in the hospital in the prior year reported lower late-life well-being ($\gamma_{05} = -4.25$) and showed steeper linear ($\gamma_{25} = -0.40$) and quadratic rates of decline ($\gamma_{45} = -0.015$). Education and material success goals were not related to individual differences in level or rates of change of well-being.

Most important for our research question, results revealed that the social orientation variables were related to individual differences in development of late-life well-being. More social participation and valuing social goals were each uniquely associated with higher well-being 1 year prior to death ($\gamma_{07} = 5.89$ and $\gamma_{08} = 2.28$,

Table 2

Single-Phase Model for Life Satisfaction Over Time to Death, Including Social Orientation and the Covariates

Parameter	Estimate	SE
Fixed effects		
Intercept, γ_{00}	45.45*	.243
Linear slope, γ_{20}	-0.90*	.045
Quadratic slope, γ_{40}	-0.03*	.002
Age at death, γ_{01}	0.09*	.015
Women, γ_{02}	-1.32*	.396
Education, γ_{03}	0.07	.098
Disability, γ_{04}	-3.46*	.390
Hospital stay last year, γ_{05}	-4.25*	.539
Material success goals, γ_{06}	0.24	.294
Social participation, γ_{07}	5.89*	.558
Social goals, γ_{08}	2.28*	.394
Family goals, γ_{09}	0.10	.243
Social Participation \times Social Goals, γ_{10}	-2.87*	.751
Change social participation, γ_{11}	-0.02	.016
Linear Slope \times Age at Death, γ_{21}	-0.003	.001
Linear Slope \times Women, γ_{22}	-0.08	.034
Linear Slope \times Education, γ_{23}	0.01	.008
Linear Slope \times Disability, γ_{24}	-0.10*	.034
Linear Slope \times Hospital Stay Last Year, γ_{25}	-0.40*	.103
Linear Slope \times Material Success Goals, γ_{26}	0.02	.026
Linear Slope \times Social Participation, γ_{27}	0.49*	.098
Linear Slope \times Social Goals, γ_{28}	0.12*	.034
Linear Slope \times Family Goals, γ_{29}	-0.05	.022
Linear Slope \times Social Participation \times Social Goals, γ_{30}	-0.14	.066
Linear Slope \times Change Social Participation, γ_{31}	0.005*	.001
Quadratic slope \times Hospital Stay Last Year, γ_{45}	-0.015**	.005
Quadratic Slope \times Social Participation, γ_{47}	0.017*	.005
Random effects		
Variance intercept	86.07*	3.275
Variance linear slope	2.17*	.135
Variance quadratic slope	0.004*	.000
Covariance Intercept, linear slope	7.10*	.550
Covariance Intercept, quadratic slope	0.19*	.025
Covariance linear slope, quadratic slope	0.09*	.006
Residual variance	64.04*	.552
Akaike information criterion	245,983	

Note. Boldface type highlights the role of social orientation for well-being late in life. Unstandardized estimates and standard errors are presented. Intercept is centered at 1 year prior to death; rate of change is scaled in T units per year as per the annual assessment schedule in the SOEP. $N = 2,910$ participants who provided 33,777 observations. Scores were standardized to a T metric ($M = 50$; $SD = 10$) using the 2002 SOEP sample as the reference frame ($M = 6.90$, $SD = 1.81$; scale range: 0–10). Two-way interaction terms of the quadratic slope and of social resource variables were tested, but only those statistically significantly different from zero were retained in the final model. SOEP = German Socio-Economic Panel Study.

* $p < .001$. ** $p = .003$.

respectively) and less steep terminal decline ($\gamma_{27} = 0.49$, $\gamma_{28} = 0.12$, and $\gamma_{48} = -0.017$, respectively). The Social Participation and Social Goals \times Level of Well-Being interaction was statistically significant and negative ($\gamma_{10} = -2.87$), indicating that the combination of compromised social participation and lack of social goals was associated with particularly low well-being at 1 year prior to death. We also found that less decline in social participation was associated with less severe rates of well-being decline ($\gamma_{31} = 0.005$). Family goals were not related to individual differences in level or rates of change of well-being.

To illustrate our major effect of interest, Figure 1 contrasts trajectories of terminal decline in well-being between participants




Figure 1. Graphic illustration of associations between levels of social orientation and late-life trajectories of well-being, as obtained from single-phase models. Participants who reported living a more socially active life (a) and those considering social goals relatively more important (b) each reported higher late-life well-being and experienced less severe terminal declines. Extrapolating from the last available information about social orientation, group differences in well-being that existed 10 years before death were exacerbated in the year of death (for the social participation groups: from 2.05 *T* score units to 6.45 *T* score units or almost two thirds of a standard deviation; for the social goals groups: from 1.91 *T* score units to 5.20 *T* score units or about half a standard deviation). Panel c illustrates the significant interaction according to which the combination of reduced social participation and lowered social goals magnified the effects of each other: The combination of compromised social participation and lack of social goals was associated with particularly low well-being at 1 year prior to death. For illustrative purposes, groups were defined using median splits. Scores for life satisfaction were standardized to a *T* metric ($M = 50$, $SD = 10$) using the 2002 German Socio-Economic Panel Study sample as the reference frame ($M = 6.90$, $SD = 1.81$; scale range: 0–10).

high and low on social participation and social goals, respectively. Using median splits for this illustration, one can see that participants who reported living a more socially active life (see Figure 1a) and those considering social goals relatively more important (see Figure 1b) each also reported higher late-life well-being and experienced less severe terminal declines. Extrapolating from the last available information about social orientation, group differences in well-being that existed 10 years before death were exacerbated in the year of death (for the social participation groups: from 2.05 *T* score units to 6.45 *T* score units or almost two thirds of a standard deviation; for the social goals groups: from 1.91 *T* score units to 5.20 *T* score units or about half a standard deviation). The significant interaction (see Figure 1c) indicates that reduced social participation and lowered social goals magnify the effects of each other. Quantifying the relations in terms of effect size, we calculated the reduction in unexplained variance in intercepts and slopes (Snijders & Bosker, 1999). Together, all predictors accounted for 20% of between-person variance in well-being levels, 7% in linear change, and 4% in quadratic change. Of these, a sizable portion was unique to the social resource variables (8.2% in levels, 3.3% in linear change, and 2.4% in quadratic change).

Does Social Orientation Relate to a Later Onset of Terminal Decline in Well-Being?

In a second set of analysis, we used two-phase models of change to examine whether social orientation and the covariates relate to

a later onset of terminal decline in well-being. Results are reported in Table 3. The typical individual transitioned into the terminal phase at $\gamma_{60} = 4.59$ years, after which the rate of decline steepened by a factor of six, from $\gamma_{20} = 0.23$ *T* score units per year to $\gamma_{40} = 1.39$ *T* score units per year. Decrements during the terminal decline phase amounted to about two thirds of a standard deviation across the last 4.5 years of life. As expected, several covariates included were associated with individual differences in levels of well-being, the onset of the terminal phase, and the rate of terminal decline. Specifically, individuals who died at older ages reported slightly higher well-being at the transition point ($\gamma_{01} = 0.13$).² Older people and women transitioned into the terminal decline phase earlier ($\gamma_{61} = -0.03$ and $\gamma_{62} = -1.01$, respectively). Participants with disability and those who had to spend one night or more in the hospital in the last year both reported lower levels of well-being at the onset of decline ($\gamma_{04} = -2.08$ and $\gamma_{05} = -2.00$, respectively), and those with hospital stays had steeper rates of terminal decline

² We probed for age differences by estimating two sets of follow-up analyses. One set excluded those who died young, operationally defined as having died before age 60 ($n = 2,503$), and another set tested for nonlinear effects by including a quadratic age term as an additional predictor. The substantive pattern of results was found to remain unchanged, and the quadratic age term did not interact with any of the social resource indicators in predicting late-life well-being and its change. We caution against overinterpreting the finding because the lack of statistically significant differences may be due to the small sample size among those who died young.

Table 3
*Multiphase Model for Life Satisfaction Over Time to Death,
 Including Social Orientation and the Covariates*

Parameter	Estimate	SE
Fixed effects		
Intercept, γ_{00}	49.58*	.204
Preterminal slope, γ_{20}	-0.23*	.020
Terminal slope, γ_{40}	-1.39*	.104
Transition point, γ_{60}	-4.59*	.017
Age at death, γ_{01}	0.13*	.012
Women, γ_{02}	-0.18	.326
Education, γ_{03}	0.05	.079
Disability, γ_{04}	-2.08*	.323
Hospital stay last year, γ_{05}	-2.00*	.406
Social participation, γ_{06}	2.17*	.404
Social goals, γ_{07}	1.04*	.315
Family goals, γ_{08}	0.53**	.200
Social Participation \times Social Goals, γ_{09}	-1.58*	.589
Change social participation, γ_{10}	-0.06*	.015
Terminal Slope \times Age at Death, γ_{41}	0.00	.005
Terminal Slope \times Women, γ_{42}	0.05	.121
Terminal Slope \times Education, γ_{43}	0.01	.032
Terminal Slope \times Disability, γ_{44}	-0.12	.1202
Terminal Slope \times Hospital Stay Last Year, γ_{45}	-0.47*	.140
Terminal Slope \times Social Participation, γ_{46}	0.03	.273
Terminal Slope \times Social Goals, γ_{47}	-0.22	.204
Terminal Slope \times Family Goals, γ_{48}	-0.15	.070
Terminal Slope \times Social Participation \times Social Goals, γ_{49}	-1.30	.522
Terminal Slope \times Change Social Participation, γ_{50}	0.00	.022
Transition Point \times Age at Death, γ_{61}	-0.03*	.002
Transition Point \times Women, γ_{62}	-1.01*	.020
Transition Point \times Education, γ_{63}	-0.01	.010
Transition Point \times Disability, γ_{64}	-0.86*	.015
Transition Point \times Hospital Stay Last Year, γ_{65}	-0.22*	.062
Transition Point \times Social Participation, γ_{66}	2.39*	.021
Transition Point \times Social Goals, γ_{67}	0.95*	.011
Transition Point \times Family Goals, γ_{68}	0.99*	.007
Transition Point \times Social Participation \times Social Goals, γ_{69}	1.71*	.018
Transition Point \times Change Social Participation, γ_{70}	0.12*	.001
Random effects		
Standard deviation intercept	8.52*	.147
Standard deviation transition point	—	—
Standard deviation preterminal slope	0.60*	.022
Standard deviation terminal slope	2.00*	.061
Correlation intercept, preterminal slope	0.54*	.026
Correlation intercept, terminal slope	-0.35*	.028
Correlation preterminal slope, preterminal slope	-0.11	.050
Residual variance	63.13*	.546
Akaike information criterion	245,778	

Note. Boldface type highlights the role of social orientation for well-being late in life. With the exception of the correlations, unstandardized estimates and standard errors are presented. Intercept centered at the transition point; rate of change scaled in T units per year as per the annual assessment schedule in the SOEP. $N = 2,910$ participants who provided 33,777 observations. Scores standardized to a T metric ($M = 50$; $SD = 10$) using the 2002 SOEP sample as the reference ($M = 6.90$, $SD = 1.81$; scale range: 0–10). SOEP = German Socio-Economic Panel Study.

* $p < .001$. ** $p = .008$.

($\gamma_{45} = -0.47$). Both disability and hospital stays in the last year were associated with spending more time in the phase of terminal decline ($\gamma_{64} = -0.86$ and $\gamma_{65} = -0.22$, respectively). Education was not related to individual differences in the two-phase trajectories of well-being.

As in the previous models, social participation and social goals were each and independently associated with higher levels of well-being at the onset of decline ($\gamma_{06} = 2.17$ and $\gamma_{07} = 1.04$, respectively) and with spending less time in the terminal decline phase ($\gamma_{66} = 2.39$ and $\gamma_{67} = 0.95$, respectively). Again, significant interaction effects emerged suggesting that the combination of high social participation and social goals being more important was associated with particularly late onset of decline in well-being ($\gamma_{69} = 1.71$). We also found that less decline in social participation was associated with a later onset of well-being decline ($\gamma_{70} = 0.17$). The multiphase model also revealed indications for the importance of family goals being associated with slightly higher levels of well-being at the onset of decline ($\gamma_{08} = 0.53$) and a later onset ($\gamma_{68} = 0.99$).

To illustrate, Figure 2 contrasts the onset of terminal decline in well-being between participants high and low on social participation and social goals, respectively. Using median splits for this illustration, one can see that participants who reported living a more socially active life (see Figure 2a) and those considering social goals relatively more important (see Figure 2b) each also reported higher late-life well-being and entered the phase of precipitous well-being decline later. The significant interaction (see Figure 2c) indicates that the combination of preserved social participation and high social goals magnifies the effects of each other, being associated with particularly late onset of decline in well-being. Figure 3 illustrates how both level and change in social participation relate to late-life trajectories of well-being. In particular, the combination of high social participation and less decline in social participation was associated with shallow rates of decline (see Figure 3a) and a late onset of decline in well-being (see Figure 3b).

The pattern of covariances also indicated that SOEP participants who reported higher well-being showed less decline in the preterminal phase ($r = .54$), but were somewhat more vulnerable to decline in the terminal phase ($r = -.35$). Interestingly, the rates of decline in the preterminal phase and the terminal phase were only marginally associated with one another ($r = -.11$, $p = .022$). Again quantifying effect size as the reduction in unexplained variance in the parameters of interest (Snijders & Bosker, 1999), we found that together all predictors accounted for 15% of between-person variance in well-being levels and 24.5% in the rate of terminal decline. Of these, a sizable portion was due to unique or shared associations of the social resource variables (8.5% in levels and 18.7% in the rate of terminal decline). Because model convergence required us to remove the random effect for the transition point, we could not use this approach to quantify effect size in the onset of decline. Using the reduction in residual variance compared with an intercept-only model as an overall effect size, the multiphase model of change and the predictors included accounted for a total of 23% of the variance in late-life trajectories of well-being. Taken together, our results suggest that levels of social goals and both levels of and changes in social activities and participation were each independently associated with the rate of late-life decline and the onset of late-life decline in well-being.

Discussion

The major objective of the current study was to move toward filling a distinct gap in the literature on potentially modifiable psychosocial individual difference factors that contribute to termi-




Figure 2. Graphic illustration of associations between levels of social orientation and late-life trajectories of well-being, as obtained from multiphase models. Participants who reported living a more socially active life (a) and those considering social goals relatively more important (b) each reported higher late-life well-being and entered the phase of precipitous well-being decline later. Panel c illustrates the significant interaction according to which the combination of social participation and social goals magnifies the effects of each other. The combination of high social participation and social goals being more important was associated with particularly late onset of decline in well-being. For illustrative purposes, groups were defined using median splits. Scores for life satisfaction were standardized to a *T* metric ($M = 50$, $SD = 10$) using the 2002 German Socio-Economic Panel Study sample as the reference frame ($M = 6.90$, $SD = 1.81$; scale range: 0–10).

nal decline in well-being. To do so, we applied single- and multiphase growth models to up to 27 years of annual longitudinal data from the SOEP. Our results extended existing evidence for the psychological benefits of social engagement into the end-of-life phase and revealed that social variables at the behavioral level (self-ratings of social participation) and at motivational levels (valuing social goals) each represent uniquely important resources for late-life well-being. Living a socially active life and prioritizing social goals were independently associated with higher late-life well-being, less severe rates of late-life declines, and a later onset of terminal decline. Significant interaction effects suggested that the combination of (reduced) social participation and (lowered) social goals magnifies the effects of each other. Findings also indicated that less decline in social participation was associated with less severe rates and a later onset of well-being decline. Interestingly, we found little evidence that valuing family goals is related to late-life trajectories of well-being. Associations were independent of key correlates of well-being and mortality, including age at death, gender, education, disability, hospital stays, and goals in other life domains. We now discuss possible pathways by which maintaining social orientation into late life may help mitigate terminal declines in well-being.

Late-Life Well-Being: The Role of Social Orientation

We are only at the very beginning of understanding the large individual differences in terminal decline in well-being (Baltes & Smith, 2003; Gerstorf & Ram, 2013). Many older adults experience a steep drop in well-being in the last years of life, whereas others maintain well-being throughout their last years. We found

that typical changes in life satisfaction at the end of life were best characterized by a multiphase model rather than a single-phase model ($\Delta = 205$ in the Akaike information criterion [AIC] for the full models and $\Delta = 77$ in the AIC for the zero-order models, see Tables 2 and 3). Nevertheless, reporting results from both models is important because the two approaches shed light on different facets of the larger phenomenon, with the single-phase models focusing on longer term trajectories of stability and change (an incremental change theory) and the multiphase models focusing on the onset of late-life decline (a stage transition theory). Though not independent of one another, the convergence of findings across approaches adds to the credibility of our results about the importance of social resources for late-life well-being. We also consider it important to keep the single-phase model because it helps connect the current study to the large majority of earlier reports on late-life well-being that were based on relatively few data points and thus could not apply multiphase models (Berg, Hassing, Thorvaldsson, & Johansson, 2011; Carmel, Shriram, & Shmotkin, 2013; Diehr, Williamson, Burke, & Psaty, 2002; Mroczek & Spiro, 2005; Palgi et al., 2010; Schilling et al., 2013; Vogel, Schilling, Wahl, Beekman, & Penninx, 2013; Windsor, Gerstorf, & Luszcz, 2015).

In the current study, we examined the role of social orientation and opted for an approach that removed differences in central background characteristics known to modulate late-life declines. Our results indicate that well-known correlates of well-being and mortality are indeed important. For example, our operational definitions of health constraints (disability and hospital stays) were associated with lower levels of late-life well-being, steeper terminal declines, and an earlier onset of decline. We found it striking




Figure 3. Graphic illustration of how level and change in social participation relate to late-life trajectories of well-being, as obtained from single-phase models (a) and multiphase models (b). In particular, the combination of high social participation and less decline in social participation was associated with less severe rates of decline and a late onset of decline in well-being. For illustrative purposes, groups were defined using median splits. Scores for life satisfaction were standardized to a T metric ($M = 50$, $SD = 10$) using the 2002 German Socio-Economic Panel Study sample as the reference frame ($M = 6.90$, $SD = 1.81$; scale range: 0–10).

to see that associations of social participation and social goals with late-life well-being were sizable and went beyond recognized sociodemographic and health factors. Setting third-variable explanations aside, results from our study can be taken to suggest that it would be all the more important for people's well-being late in life if they could continue to set social goals and carry on with social activities despite the often pervasive challenges they confronted. To illustrate, compromised well-being late in life often results from the loss of social belonging (e.g., as a consequence of widowhood and other social losses) and the increased difficulties with realizing one's social goals (e.g., as a consequence of poor functional health; Gerstorf & Ram, 2015; Rook, Mavandadi, Sorokin, & Zettel, 2007; Windsor et al., 2015). Future, more mechanism-oriented research would need to thoroughly disentangle key questions about the temporal and causal ordering.

Social activities and participation. In line with reports from younger samples (Huxhold et al., 2013), our results suggest that continuing to participate in social activities in the last years is associated with maintaining well-being. Investing one's physical and psychological resources into socially oriented activities of a sharing or instrumental kind can be expected to be advantageous at a number of different levels. Well-being may be boosted directly by carrying out joyful activities (Lyubomirsky & Layous, 2013) or

indirectly by facilitating self-esteem and a sense of control or promote physical and cognitive functioning (Leary & Baumeister, 2000; Zautra, Infurna, Zautra, Gallardo, & Velasco, 2016). For example, Lövdén, Ghisletta, and Lindenberger (2005) reported that social participation precedes and predicts less severe rates of decline in perceptual speed as a measure of cognitive functioning. The authors argued that an engaged lifestyle often involves cognitive stimulation and physical activity, which in turn protect against neurophysiological and cardiovascular pathology underlying cognitive decline. By virtue of maintaining cognitive functioning into late life, it may also be easier to maintain well-being.

The importance of social and family goals. Despite differences in goal contents, goals are highly important in shaping well-being and other mental health outcomes across adulthood because they form the basis for subjective evaluations of whether a person is moving in a desired direction (Austin & Vancouver, 1996; Nurmi & Salmela-Aro, 2006). Notably, goals not only contribute to well-being, but they can also reveal constraints that draw attention to the importance of strategies to make the best out of limited resources. SST as one of the major theories in the field proposes that recognition that future time is limited rather than open-ended prompts motivational shifts that lead individuals to prioritize socioemotional goals and prefer familiar social partners,

which in turn helps maintain well-being (Carstensen, 2006; Carstensen et al., 2011). We acknowledge that our data did not allow for contrasting these social goals with knowledge acquisition goals. Nevertheless, our results extend these notions to the very end of life and are consistent with the idea that prioritizing social goals at the end of life is associated with maintaining well-being as long as possible. Valuing and pursuing social goals may contribute to well-being by boosting feelings of competence, generativity, and belonging.

It was beyond the scope and possibilities of our study to examine the specific mechanisms by which social orientation operates to help people maintain well-being at the end of life. However, insights into how social factors operate may help to design targeted interventions aimed at reshaping the social context in order to assuage terminal decline in well-being. In this respect, it is particularly interesting that the combination of compromised social participation and low endorsement of social goals appeared to magnify the effects of each other. One possible interpretation would be that social deprivation combined with giving up on social goals may be particularly vicious because it not only eliminates an important source of well-being concurrently, but it also reflects the individual's perception that this is something that he or she cannot change in the future. It is also possible that participants in this group may have experienced demoralization syndrome (Robinson, Kissane, Brooker, & Burney, 2015), a state of existential distress in the face of progressive disease that is characterized by helplessness and hopelessness, lack of goals and meaning, high levels of negative emotions, and profound social isolation. The growing literature on this condition, which is estimated to affect up to a third of individuals with progressive medical conditions, may ultimately point to promising pathways for intervention.

Participants in our study were also asked to rate the importance of having a good marriage and a good relationship with their children (i.e., family-related goals). Consistent with reports from earlier phases of life (Lang, 2001; Neyer & Lang, 2003; Silverstein & Bengtson, 1994), we had expected that having good relationships with one's close family members would be central for people's well-being at the end of life. For example, if people felt that their children had turned out well and they had good relationships with them, then this would contribute to a sense of generativity (Erikson, 1985). It may be easier to let go and accept things the way they are when people have the impression that they have left a lasting legacy in this world. It was thus surprising that prioritizing family goals evinced very few statistically significant associations with well-being after social participation and the importance of social goals were taken into account. We note that even fewer associations existed at the zero-order level when examined as a separate predictor. We can only speculate about possible reasons. One interpretation is that the response distribution was too skewed, with an average of 3.15 on a 1-to-4 response scale for the family goal items compared with an average of 2.33 for the social goals (see Table 1). This may in part reflect a social desirability bias. Most important, however, the item and response format for family goals may not have been fine-grained enough to capture subtle, but salient individual differences in valuing family goals that distinguish between those who declined and those who maintained well-being late in life. A second interpretation is that participants who had lost their partners, never married, and/or never had (or had lost their) children were, of course, unable to

answer questions about the importance of spouses and/or children. It is thus possible that earlier obtained data (i.e., further away from death) were used for those who experienced late life without partner or child, whereas later obtained data (i.e., closer to death) were used for those who experienced late life with their partner and or child. If this confound exists, it may have contributed to the lack of differentiation we had observed for the variable indicating a prioritization of partner and children. Empirically, however, it is not the case that the time elapsed between goal assessments and death was larger for family goals ($M = 4.11$ years, $SD = 3.38$) than for goals in the other two domains of life (material success: $M = 4.08$ years, $SD = 3.35$; social: $M = 4.08$ years, $SD = 3.35$). The probably more general point for future inquiry would be to examine what adaptive goal structures in late life look like for people for whom engagement with family is not an option (any-more).

Apart from such methodological concerns, it is possible that our findings accurately reflect the complexity of family relationships in later life. Family life is often a "mixed bag" and represents a source of not only joy, but also worry and tensions, stress and sorrow (Birditt, Fingerman, & Almeida, 2005; Hoppmann & Gerstorf, 2016). For example, valuing one's partner often makes people vulnerable to declines in well-being when the partner suffers from cognitive or physical limitations (Michalowski, Hoppmann, & Gerstorf, 2014) and relationships with both children and spouses are further complicated with the onset of caregiving responsibilities (Zarit & Reamy, 2013). In a similar vein, relationships with adult children can be ambivalent even for healthy older adults, especially when children differ in values and have not attained educational and interpersonal success (Fingerman, Chen, Hay, Cichy, & Lefkowitz, 2006; Pillemer, Munsch, Fuller-Rowell, Riffin, & Suitor, 2012). To disentangle this complexity, future research should differentiate between goals for adult children and spouses and control for caregiving responsibilities and relationship quality.

Limitations and Outlook

The larger sample size available in the current study and advancements in modeling efforts allowed for a more thorough test of our research questions than has been previously possible. For example, we were able to estimate models that simultaneously included associations of the social resource variables and of the covariates with all three aspects of late-life trajectories in which we are primarily interested (i.e., level, rates of terminal decline, and onset of decline). This was not previously possible. For example, the 1,641 participants in Gerstorf et al. (2014) only allowed for estimating associations of the predictor variables with the onset of decline, but not parallel associations with level and the rate of change.

Despite these advances, there are important limitations to our study. To begin, we acknowledge that many pivotal questions related to our results remain open, particularly those that involve the mechanisms underlying the role of social orientation for terminal decline in well-being. If terminal decline (in well-being) reflects some combination of late-life neuropathology (e.g., Alzheimer disease, Lewy bodies), deteriorating integrity of neurocognitive control systems, and a breakdown of overall system coordination and integrity, then key questions are which social features

(e.g., instrumental vs. emotional) operate in what ways (e.g., direct vs. indirect) to alleviate the consequences of these processes. In a similar vein, future research may benefit from disentangling whether effects observed for social participation are driven primarily by the social nature of the activity or the preserved physical fitness that is needed to lead a socially active life (Berkman et al., 2000). Even if the underlying mechanisms can be established, one of the next questions is how social indicators of quality of life can be maintained or reestablished late in life.

Second, social orientation and social resources constitute a broad construct space. In the current study, we focused on process aspects of the Fingerman and Lang (2004) framework of social relations, but did not investigate specifically how facets of structure (e.g., parent-child, romantic partner, sibling, friend) or outcome operate (e.g., social support, relationship quality). Specifically, we used two distinct social orientations that tap into largely different aspects of that construct space, namely, the social activities people pursue and the social and family goals they have and value. However, measures about further highly informative aspects of the larger construct space are not included in the SOEP assessment battery, and have thus not been available for our secondary data analysis. For example, loneliness, feeling part of a group, receiving instrumental and emotional support from others, or perceived available resources such as number and perceived quality of active social ties can each be expected to play a role for late-life well-being. To illustrate, measures of social support would have allowed exploring how people at the end of life continue to actively seek out, mobilize, and handle the availability of social support (Lang, Featherman, & Nesselrode, 1997). Beneficial features of social support include encouraging mastery attempts, providing supportive feedback, and suggesting appropriate coping responses in times of strain. Combining the results of the current study with recent reports about the importance of perceived personal control late in life (Gerstorf et al., 2014; Heckhausen et al., 2013), it would be particularly intriguing to examine how social orientation and social resources interact with (e.g., boost) perceived personal control so to maintain well-being and health as long as possible (Antonucci, 2001; Cohen & Wills, 1985). Because measures of social orientation and those of personal control were assessed at different points in time over the 30 years the SOEP study has now run, it was unfortunately not an option to examine social resources and personal control simultaneously in our analyses.

Third, it would have been feasible to directly model how both social orientation and well-being develop in parallel at the end of life, to test other than linear approximations of change for the social resource variables, and to examine whether well-being operates as a driving force for changes in social orientation. For example, it is possible that compromised well-being sets a downward spiral in action that prompts people to live a more and more restricted social life (e.g., because social activities are increasingly devalued; Achterberg et al., 2003). However, limitations of our data set did not allow us to test these intriguing questions. In particular, data for the social resources were available for considerably fewer measurement occasions (e.g., no more than five waves for social goals) than for well-being (up to 27 waves). As a consequence, our modeling efforts already pushed the envelope of what is currently possible with the data at hand. We thus chose to prioritize the robustness of the models, which would have been

compromised by extensions to single-step multivariate (parallel process) models. In addition, we estimated follow-up analyses that either also included quadratic rates of change for social participation or modeled change in social orientation over a time-to-death metric. The substantive pattern of results remained unchanged in these follow-up analyses (e.g., less decline in social participation with impending death was associated with less severe rates and a later onset of well-being decline). We hesitate to present the time-to-death modeling of the social orientation variables as our main analyses because the scarce assessment schedules relative to the well-being measure would not permit considering and modeling these as parallel processes. In our view, modeling the social orientation variables with linear change over time provides a more conservative estimate and does not convey the impression of overselling our design. Our cautious interpretation and handling of changes in social orientation in our study is further supported empirically by the average changes obtained, which was modest in size: yearly change on social participation, -0.005 on a 1–4 scale ($p = .0011$); on social goals, -0.015 on a 1–4 scale ($p = .0003$); and family goals, 0.002 on a 1–4 scale ($p = .76$), with no evidence for statistically significant quadratic changes.

Finally, the SOEP is a great resource that has tracked people annually for almost 30 years now, thereby having provided considerable insights into the stability and change of central domains of functioning across adulthood. However, it does not provide information about cause of death. Thus, we could not screen out accidental and other sudden causes of death not preceded by a prodromal period of decline, nor could we differentiate dementias from other chronic conditions. It is possible that social resources are particularly important for late-life well-being when people are confronted with debilitating chronic health conditions (Hoppmann & Gerstorf, 2016). Furthermore, the annual assessment schedule approaches its limits when it comes to better understanding the very last months of life. Very little information is available about well-being and the social resources in the last 12 months of life. The current study thus sheds light on the terminal well-being changes people experience and the resources people bring into the last year of life, but we do not know how preserved or impaired well-being and the (social) resources were in the very last months of life. Similarly, information about how people value certain life goals was obtained an average of 2–4 years before death. With a more frequent sampling scheme zooming in on the last months, we would expect rapid average well-being declines (using proxy reports; see Infurna et al., 2014) and probably a more succinct pattern for the role of social orientation and social resources. A hypothesis to test once appropriate data are available would be whether people maintain reasonably high well-being if they can preserve their social orientation, receive the instrumental and emotional support needed, and are able to continue pursuing a minimum amount of social activities. Finally, our selectivity analyses conjointly suggest that SOEP participants who died at an older age, received less education, and started with lower levels of well-being were less likely to remain in our study until close to death and provided fewer assessments of life satisfaction. As a consequence, the results we obtained probably do not generalize to less positively selected population segments than those included in our report. It would also be intriguing to more closely examine possible cultural differences in how effective social resources may

operate (e.g., family-focused networks being as beneficial as diverse networks in China; Cheng, Lee, Chan, Leung, & Lee, 2009).

Conclusion

The current study is one of very few reports to identify psychosocial characteristics associated with vulnerability to or resilience in the face of terminal decline in well-being. Taking into account central covariates of well-being and mortality, our results indicate that social orientation is related to maintaining well-being as long as possible into the very last years of life. Building on and extending work on earlier phases of life, our results suggest that older adults, even at the end of life, continue to play an active role in shaping their own well-being by engaging in social activities and by prioritizing social goals. Much more needs to be learned, however, about the underlying mechanisms.

References

- Achterberg, W., Pot, A. M., Kerkstra, A., Ooms, M., Muller, M., & Ribbe, M. (2003). The effect of depression on social engagement in newly admitted Dutch nursing home residents. *The Gerontologist, 43*, 213–218. <http://dx.doi.org/10.1093/geront/43.2.213>
- Antonucci, T. C. (2001). Social relations: An examination of social networks, social support, and sense of control. In J. E. Birren & K. W. Schaie (Eds.), *Handbook of the psychology of aging* (pp. 427–453). San Diego, CA: Academic Press.
- Atchley, R. C. (1989). A continuity theory of normal aging. *The Gerontologist, 29*, 183–190. <http://dx.doi.org/10.1093/geront/29.2.183>
- Austin, J. T., & Vancouver, J. B. (1996). Goal constructs in psychology: Structure, process, and content. *Psychological Bulletin, 120*, 338–375. <http://dx.doi.org/10.1037/0033-2909.120.3.338>
- Baltes, P., & Nesselroade, J. (1979). History and rationale of longitudinal research. In J. R. Nesselroade & P. Baltes (Eds.), *Longitudinal research in the study of behavior and development* (pp. 1–39). San Diego, CA: Academic Press.
- Baltes, P. B., & Smith, J. (2003). New frontiers in the future of aging: From successful aging of the young old to the dilemmas of the fourth age. *Gerontology, 49*, 123–135. <http://dx.doi.org/10.1159/000067946>
- Berg, A. I., Hassing, L. B., Thorvaldsson, V., & Johansson, B. (2011). Personality and personal control make a difference for life satisfaction in the oldest-old: Findings in a longitudinal population-based study of individuals 80 and older. *European Journal of Ageing, 8*, 13–20. <http://dx.doi.org/10.1007/s10433-011-0181-9>
- Berkman, L. F., Glass, T., Brissette, I., & Seeman, T. E. (2000). From social integration to health: Durkheim in the new millennium. *Social Science & Medicine, 51*, 843–857. [http://dx.doi.org/10.1016/S0277-9536\(00\)00065-4](http://dx.doi.org/10.1016/S0277-9536(00)00065-4)
- Birditt, K. S., Fingerman, K. L., & Almeida, D. M. (2005). Age differences in exposure and reactions to interpersonal tensions: A daily diary study. *Psychology and Aging, 20*, 330–340. <http://dx.doi.org/10.1037/0882-7974.20.2.330>
- Brockmann, H., & Klein, T. (2004). Love and death in Germany: The marital biography and its effect on mortality. *Journal of Marriage and Family, 66*, 567–581. <http://dx.doi.org/10.1111/j.0022-2445.2004.00038.x>
- Burgess, E. W. (1954). Social relations, activities, and personal adjustment. *American Journal of Sociology, 59*, 352–360. <http://dx.doi.org/10.1086/221368>
- Burkhauser, R. V., Giles, P., Lillard, D. R., & Schwarze, J. (2005). Until death do us part: An analysis of the economic well-being of widows in four countries. *The Journals of Gerontology. Series B: Psychological Sciences and Social Sciences, 60*, S238–S246. <http://dx.doi.org/10.1093/geronb/60.5.S238>
- Burns, R. A., Byles, J., Magliano, D. J., Mitchell, P., & Anstey, K. J. (2015). The utility of estimating population-level trajectories of terminal wellbeing decline within a growth mixture modeling framework. *Social Psychiatry and Psychiatric Epidemiology, 50*, 479–487. <http://dx.doi.org/10.1007/s00127-014-0948-3>
- Burns, R. A., Mitchell, P., Shaw, J., & Anstey, K. J. (2014). Trajectories of terminal decline in the well-being of older women: The DYNOPTA project. *Psychology and Aging, 29*, 44–56. <http://dx.doi.org/10.1037/a0035370>
- Carmel, S., Shrira, A., & Shmotkin, D. (2013). The will to live and death-related decline in life satisfaction. *Psychology and Aging, 28*, 1115–1123. <http://dx.doi.org/10.1037/a0034649>
- Carstensen, L. L. (1995). Evidence for a life-span theory of socioemotional selectivity. *Current Directions in Psychological Science, 4*, 151–156. <http://dx.doi.org/10.1111/1467-8721.ep11512261>
- Carstensen, L. L. (2006). The influence of a sense of time on human development. *Science, 312*, 1913–1915. <http://dx.doi.org/10.1126/science.1127488>
- Carstensen, L. L., Turan, B., Scheibe, S., Ram, N., Ersner-Hershfield, H., Samanez-Larkin, G. R., . . . Nesselroade, J. R. (2011). Emotional experience improves with age: Evidence based on over 10 years of experience sampling. *Psychology and Aging, 26*, 21–33. <http://dx.doi.org/10.1037/a0021285>
- Charles, S. T. (2010). Strength and vulnerability integration: A model of emotional well-being across adulthood. *Psychological Bulletin, 136*, 1068–1091. <http://dx.doi.org/10.1037/a0021232>
- Charles, S. T., Reynolds, C. A., & Gatz, M. (2001). Age-related differences and change in positive and negative affect over 23 years. *Journal of Personality and Social Psychology, 80*, 136–151. <http://dx.doi.org/10.1037/0022-3514.80.1.136>
- Cheng, S.-T., Lee, C. K. L., Chan, A. C. M., Leung, E. M. F., & Lee, J.-J. (2009). Social network types and subjective well-being in Chinese older adults. *The Journals of Gerontology. Series B: Psychological Sciences and Social Sciences, 64B*, 713–722. <http://dx.doi.org/10.1093/geronb/gbp075>
- Cohen, S. (2004). Social relationships and health. *American Psychologist, 59*, 676–684. <http://dx.doi.org/10.1037/0003-066X.59.8.676>
- Cohen, S., & Wills, T. A. (1985). Stress, social support, and the buffering hypothesis. *Psychological Bulletin, 98*, 310–357. <http://dx.doi.org/10.1037/0033-2909.98.2.310>
- Cudeck, R., & Harring, J. R. (2007). Analysis of nonlinear patterns of change with random coefficient models. *Annual Review of Psychology, 58*, 615–637. <http://dx.doi.org/10.1146/annurev.psych.58.110405.085520>
- Cudeck, R., & Klebe, K. J. (2002). Multiphase mixed-effects models for repeated measures data. *Psychological Methods, 7*, 41–63. <http://dx.doi.org/10.1037/1082-989X.7.1.41>
- Dahlberg, L., & McKee, K. J. (2014). Correlates of social and emotional loneliness in older people: Evidence from an English community study. *Aging & Mental Health, 18*, 504–514. <http://dx.doi.org/10.1080/13607863.2013.856863>
- DeNeve, K. M., & Cooper, H. (1998). The happy personality: A meta-analysis of 137 personality traits and subjective well-being. *Psychological Bulletin, 124*, 197–229. <http://dx.doi.org/10.1037/0033-2909.124.2.197>
- Diehr, P., Williamson, J., Burke, G. L., & Psaty, B. M. (2002). The aging and dying processes and the health of older adults. *Journal of Clinical Epidemiology, 55*, 269–278. [http://dx.doi.org/10.1016/S0895-4356\(01\)00462-0](http://dx.doi.org/10.1016/S0895-4356(01)00462-0)
- Diener, E., Lucas, R. E., & Scollon, C. N. (2006). Beyond the hedonic treadmill: Revising the adaptation theory of well-being. *American Psychologist, 61*, 305–314. <http://dx.doi.org/10.1037/0003-066X.61.4.305>

- Diener, E., & Seligman, M. E. P. (2004). Beyond money: Toward an economy of well-being. *Psychological Science in the Public Interest*, 5, 1–31. <http://dx.doi.org/10.1111/j.0963-7214.2004.00501001.x>
- Erikson, E. H. (1985). *The life cycle completed: A review*. New York, NY: Norton.
- Fingerman, K. L., Chen, P.-C., Hay, E., Cichy, K. E., & Lefkowitz, E. S. (2006). Ambivalent reactions in the parent and offspring relationship. *The Journals of Gerontology*. Series B: Psychological Sciences and Social Sciences, 61, P152–P160. <http://dx.doi.org/10.1093/geronb/61.3.P152>
- Fingerman, K. L., & Lang, F. R. (2004). Coming together: A perspective on relationships across the life span. In F. R. Lang & K. L. Fingerman (Eds.), *Growing together: Personal relationships across the life span* (pp. 1–13). New York, NY: Cambridge University Press.
- Fredrickson, B. L., & Carstensen, L. L. (1990). Choosing social partners: How old age and anticipated endings make people more selective. *Psychology and Aging*, 5, 335–347. <http://dx.doi.org/10.1037/0882-7974.5.3.335>
- Fujita, F., & Diener, E. (2005). Life satisfaction set point: Stability and change. *Journal of Personality and Social Psychology*, 88, 158–164.
- Fung, H. H., Carstensen, L. L., & Lutz, A. M. (1999). Influence of time on social preferences: Implications for life-span development. *Psychology and Aging*, 14, 595–604. <http://dx.doi.org/10.1037/0882-7974.14.4.595>
- Gerstorf, D., Heckhausen, J., Ram, N., Infurna, F. J., Schupp, J., & Wagner, G. G. (2014). Perceived personal control buffers terminal decline in well-being. *Psychology and Aging*, 29, 612–625. <http://dx.doi.org/10.1037/a0037227>
- Gerstorf, D., & Ram, N. (2013). Inquiry into terminal decline: Five objectives for future study. *The Gerontologist*, 53, 727–737. <http://dx.doi.org/10.1093/geront/gnt046>
- Gerstorf, D., & Ram, N. (2015). A framework for studying mechanisms underlying terminal decline in well-being. *International Journal of Behavioral Development*, 39, 210–220. <http://dx.doi.org/10.1177/0165025414565408>
- Gerstorf, D., Ram, N., Lindenberger, U., & Smith, J. (2013). Age and time-to-death trajectories of change in indicators of cognitive, sensory, physical, health, social, and self-related functions. *Developmental Psychology*, 49, 1805–1821. <http://dx.doi.org/10.1037/a0031340>
- Gerstorf, D., Ram, N., Mayraz, G., Hidajat, M., Lindenberger, U., Wagner, G. G., & Schupp, J. (2010). Late-life decline in well-being across adulthood in Germany, the United Kingdom, and the United States: Something is seriously wrong at the end of life. *Psychology and Aging*, 25, 477–485. <http://dx.doi.org/10.1037/a0017543>
- Glass, T. A., Seeman, T. E., Herzog, A. R., Kahn, R., & Berkman, L. F. (1995). Change in productive activity in late adulthood: MacArthur Studies of Successful Aging. *The Journals of Gerontology*. Series B: Psychological Sciences and Social Sciences, 50B, S65–S76. <http://dx.doi.org/10.1093/geronb/50B.2.S65>
- Guralnik, J. M. (1991). Prospects for the compression of morbidity: The challenge posed by increasing disability in the years prior to death. *Journal of Aging and Health*, 3, 138–154. <http://dx.doi.org/10.1177/089826439100300202>
- Hawton, A., Green, C., Dickens, A. P., Richards, S. H., Taylor, R. S., Edwards, R., . . . Campbell, J. L. (2011). The impact of social isolation on the health status and health-related quality of life of older people. *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation*, 20, 57–67. <http://dx.doi.org/10.1007/s11136-010-9717-2>
- Headey, B. (2008). Life goals matter to happiness: A revision of set-point theory. *Social Indicators Research*, 86, 213–231. <http://dx.doi.org/10.1007/s11205-007-9138-y>
- Headey, B., Muffels, R., & Wagner, G. G. (2010). Long-running German panel survey shows that personal and economic choices, not just genes, matter for happiness. *PNAS: Proceedings of the National Academy of Sciences of the United States of America*, 107, 17922–17926. <http://dx.doi.org/10.1073/pnas.1008612107>
- Heckhausen, J., Wrosch, C., & Schulz, R. (2010). A motivational theory of life-span development. *Psychological Review*, 117, 32–60. <http://dx.doi.org/10.1037/a0017668>
- Heckhausen, J., Wrosch, C., & Schulz, R. (2013). A lines-of-defense model for managing health threats: A review. *Gerontology*, 59, 438–447. <http://dx.doi.org/10.1159/000351269>
- Herzog, A. R., Franks, M. M., Markus, H. R., & Holmberg, D. (1998). Activities and well-being in older age: Effects of self-concept and educational attainment. *Psychology and Aging*, 13, 179–185. <http://dx.doi.org/10.1037/0882-7974.13.2.179>
- Herzog, A. R., & House, J. S. (1991). Productive activities and aging well. *Generations*, 15, 49–54.
- Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine*, 7, e1000316. <http://dx.doi.org/10.1371/journal.pmed.1000316>
- Hoppmann, C. A., & Gerstorf, D. (2016). Social interrelations in aging: The sample case of married couples. In K. W. Schaie & S. L. Willis (Eds.), *Handbook of the psychology of aging* (8th ed., pp. 263–277). San Diego, CA: Elsevier. <http://dx.doi.org/10.1016/B978-0-12-411469-2.00014-5>
- Hoppmann, C. A., Gerstorf, D., Smith, J., & Klumb, P. L. (2007). Linking possible selves and behavior: Do domain-specific hopes and fears translate into daily activities in very old age? *The Journals of Gerontology*. Series B: Psychological Sciences and Social Sciences, 62, P104–P111. <http://dx.doi.org/10.1093/geronb/62.2.P104>
- Huxhold, O., Fiori, K. L., & Windsor, T. D. (2013). The dynamic interplay of social network characteristics, subjective well-being, and health: The costs and benefits of socio-emotional selectivity. *Psychology and Aging*, 28, 3–16. <http://dx.doi.org/10.1037/a0030170>
- Infurna, F. J., Gerstorf, D., Ram, N., Schupp, J., Sprangers, M. A. G., & Wagner, G. G. (2014). Linking concurrent self-reports and retrospective proxy reports about the last year of life: A prevailing picture of life satisfaction decline. *The Journals of Gerontology*. Series B: Psychological Sciences and Social Sciences, 69, 695–709. <http://dx.doi.org/10.1093/geronb/gbt055>
- Infurna, F. J., Gerstorf, D., Ram, N., Schupp, J., & Wagner, G. G. (2011). Long-term antecedents and outcomes of perceived control. *Psychology and Aging*, 26, 559–575. <http://dx.doi.org/10.1037/a0022890>
- Klein, T. (1996). Determinants of institutionalization in old age. In R. Eisen & F. Sloan (Eds.), *Long-term care: Economic issues and policy solutions* (pp. 103–113). Boston, MA: Kluwer. http://dx.doi.org/10.1007/978-1-4615-4096-0_6
- Kroh, M. (2014). *Documentation of sample sizes and panel attrition in the German Socio Economic Panel, SOEP (1984 until 2012)* (SOEP Survey Papers 177: Series D). Berlin, Germany: DIW/SOEP.
- Kroh, M., Pischner, R., Spieß, M., & Wagner, G. G. (2008). On the treatment of non-original sample members in the German Household Panel Study (SOEP): Tracing and weighting. *Methoden–Daten–Analysen*, 2, 179–198.
- Lang, F. R. (2000). Endings and continuity of social relationships: Maximizing intrinsic benefits within personal networks when feeling near to death. *Journal of Social and Personal Relationships*, 17, 155–182. <http://dx.doi.org/10.1177/0265407500172001>
- Lang, F. R. (2001). Regulation of social relationships in later adulthood. *The Journals of Gerontology*. Series B: Psychological Sciences and Social Sciences, 56, P321–P326. <http://dx.doi.org/10.1093/geronb/56.6.P321>
- Lang, F. R., & Carstensen, L. L. (1994). Close emotional relationships in late life: Further support for proactive aging in the social domain. *Psychology and Aging*, 9, 315–324. <http://dx.doi.org/10.1037/0882-7974.9.2.315>

- Lang, F. R., & Carstensen, L. L. (2002). Time counts: Future time perspective, goals, and social relationships. *Psychology and Aging, 17*, 125–139. <http://dx.doi.org/10.1037/0882-7974.17.1.125>
- Lang, F. R., Featherman, D. L., & Nesselrode, J. R. (1997). Social self-efficacy and short-term variability in social relationships: The MacArthur Successful Aging Studies. *Psychology and Aging, 12*, 657–666. <http://dx.doi.org/10.1037/0882-7974.12.4.657>
- Lawton, M. P., Ruckdeschel, K., Winter, L., & Kleban, M. H. (1999). Affect-experiential personality types in middle and late adulthood. *Journal of Mental Health and Aging, 5*, 223–239.
- Leary, M. R., & Baumeister, R. F. (2000). The nature and function of self-esteem: Sociometer theory. *Advances in Experimental Social Psychology, 32*, 1–62. [http://dx.doi.org/10.1016/S0065-2601\(00\)80003-9](http://dx.doi.org/10.1016/S0065-2601(00)80003-9)
- Lemon, B. W., Bengtson, V. L., & Peterson, J. A. (1972). An exploration of the activity theory of aging: Activity types and life satisfaction among in-movers to a retirement community. *The Journals of Gerontology, 27*, 511–523. <http://dx.doi.org/10.1093/geronj/27.4.511>
- Littell, R. C., Milliken, G. A., Stroup, W. W., Wolfinger, R. D., & Schabenberger, O. (2006). *SAS for mixed models* (2nd ed.). Cary, NC: SAS Institute.
- Little, R. J. A., & Rubin, D. B. (1987). *Statistical analysis with missing data*. New York, NY: Wiley.
- Litwin, H., & Shiovitz-Ezra, S. (2006). The association between activity and wellbeing in later life: What really matters? *Ageing & Society, 26*, 225–242. <http://dx.doi.org/10.1017/S0144686X05004538>
- Litwin, H., & Stoeckel, K. J. (2014). Confidant network types and well-being among older Europeans. *The Gerontologist, 54*, 762–772. <http://dx.doi.org/10.1093/geront/gnt056>
- Lövdén, M., Ghisletta, P., & Lindenberger, U. (2005). Social participation attenuates decline in perceptual speed in old and very old age. *Psychology and Aging, 20*, 423–434. <http://dx.doi.org/10.1037/0882-7974.20.3.423>
- Lucas, R. E. (2007). Long-term disability is associated with lasting changes in subjective well-being: Evidence from two nationally representative longitudinal studies. *Journal of Personality and Social Psychology, 92*, 717–730. <http://dx.doi.org/10.1037/0022-3514.92.4.717>
- Lucas, R. E., Clark, A. E., Georgellis, Y., & Diener, E. (2003). Reexamining adaptation and the set point model of happiness: Reactions to changes in marital status. *Journal of Personality and Social Psychology, 84*, 527–539. <http://dx.doi.org/10.1037/0022-3514.84.3.527>
- Lyubomirsky, S., & Layous, K. (2013). How do simple positive activities increase well-being? *Current Directions in Psychological Science, 22*, 57–62. <http://dx.doi.org/10.1177/0963721412469809>
- McArdle, J. J. (1994). Structural factor analysis experiments with incomplete data. *Multivariate Behavioral Research, 29*, 409–454. http://dx.doi.org/10.1207/s15327906mbr2904_5
- McArdle, J. J., & Nesselrode, J. R. (2003). Growth curve analysis in contemporary psychological research. In J. Schinka & W. Velicer (Eds.), *Comprehensive handbook of psychology: Vol. 2. Research methods in psychology* (pp. 447–480). New York, NY: Pergamon Press. <http://dx.doi.org/10.1002/0471264385.wei0218>
- Menec, V. H. (2003). The relation between everyday activities and successful aging: A 6-year longitudinal study. *The Journals of Gerontology. Series B: Psychological Sciences and Social Sciences, 58*, S74–S82. <http://dx.doi.org/10.1093/geronb/58.2.S74>
- Michalowski, V. I., Hoppmann, C. A., & Gerstorf, D. (2014). Associations between perceived support in older adult marriages and dyadic covariations in momentary affect and aches. *Journals of Gerontology, Series B: Psychological Sciences and Social Sciences*. <http://dx.doi.org/10.1093/geronb/gbu151>
- Moen, P. (2001). The gendered life course. In R. H. Binstock & L. K. George (Eds.), *Handbook of aging and the social sciences* (Vol. 5, pp. 179–196). San Diego, CA: Academic Press.
- Mroczek, D. K., & Spiro, A., III. (2005). Change in life satisfaction during adulthood: Findings from the Veterans Affairs Normative Aging Study. *Journal of Personality and Social Psychology, 88*, 189–202.
- Mutran, E. (1987). Family, social ties and self-meaning in old age: The Development of an affective identity. *Journal of Social and Personal Relationships, 4*, 463–480. <http://dx.doi.org/10.1177/0265407587044005>
- Netz, Y., Wu, M. J., Becker, B. J., & Tenenbaum, G. (2005). Physical activity and psychological well-being in advanced age: A meta-analysis of intervention studies. *Psychology and Aging, 20*, 272–284. <http://dx.doi.org/10.1037/0882-7974.20.2.272>
- Neyer, F. J., & Lang, F. R. (2003). Blood is thicker than water: Kinship orientation across adulthood. *Journal of Personality and Social Psychology, 84*, 310–321. <http://dx.doi.org/10.1037/0022-3514.84.2.310>
- Nurmi, J.-E., & Salmela-Aro, K. (2006). What works makes you happy: The role of personal goals in life-span development. In M. Csikszentmihalyi & I. S. Csikszentmihalyi (Eds.), *A life worth living: Contributions to positive psychology* (pp. 182–199). New York, NY: Oxford University Press.
- Ong, A. D., Rothstein, J. D., & Uchino, B. N. (2012). Loneliness accentuates age differences in cardiovascular responses to social evaluative threat. *Psychology and Aging, 27*, 190–198. <http://dx.doi.org/10.1037/a0025570>
- Palgi, Y., Shrira, A., Ben-Ezra, M., Spalter, T., Shmotkin, D., & Kavé, G. (2010). Delineating terminal change in subjective well-being and subjective health. *The Journals of Gerontology. Series B: Psychological Sciences and Social Sciences, 65B*, 61–64. <http://dx.doi.org/10.1093/geronb/gbp095>
- Parslow, R. A., Jorm, A. F., Christensen, H., & Mackinnon, A. (2006). An instrument to measure engagement in life: Factor analysis and associations with sociodemographic, health and cognition measures. *Gerontology, 52*, 188–198. <http://dx.doi.org/10.1159/000091829>
- Pillemer, K., Munsch, C. L., Fuller-Rowell, T., Riffin, C., & Sutor, J. J. (2012). Ambivalence toward adult children: Differences between mothers and fathers. *Journal of Marriage and Family, 74*, 1101–1113. <http://dx.doi.org/10.1111/j.1741-3737.2012.01004.x>
- Prince-Paul, M. (2008). Understanding the meaning of social well-being at the end of life. *Oncology Nursing Forum, 35*, 365–371. <http://dx.doi.org/10.1188/08.ONF.365-371>
- Ram, N., & Grimm, K. (2015). Growth curve modeling and longitudinal factor analysis. In W. F. Overton & P. C. M. Molenaar (Eds.), *Handbook of child psychology and developmental science: Vol. 1. Theory and method* (7th ed., pp. 758–788). Hoboken, NJ: Wiley. <http://dx.doi.org/10.1002/9781118963418.childpsy120>
- Rehse, B., & Pukrop, R. (2003). Effects of psychosocial interventions on quality of life in adult cancer patients: Meta analysis of 37 published controlled outcome studies. *Patient Education and Counseling, 50*, 179–186. [http://dx.doi.org/10.1016/S0738-3991\(02\)00149-0](http://dx.doi.org/10.1016/S0738-3991(02)00149-0)
- Robinson, S., Kissane, D. W., Brooker, J., & Burney, S. (2015). A systematic review of the demoralization syndrome in individuals with progressive disease and cancer: A decade of research. *Journal of Pain and Symptom Management, 49*, 595–610. <http://dx.doi.org/10.1016/j.jpainsymman.2014.07.008>
- Rook, K. S., Mavandadi, S., Sorkin, D. H., & Zettel, L. A. (2007). Optimizing social relationships as a resource for health and well-being in later life. In C. M. Aldwin, C. L. Park, & A. Spiro (Eds.), *Handbook of health psychology and aging* (pp. 267–285). New York, NY: Guilford Press.
- Rowe, J. W., & Kahn, R. L. (1997). Successful aging. *The Gerontologist, 37*, 433–440. <http://dx.doi.org/10.1093/geront/37.4.433>
- Rowe, J. W., & Kahn, R. L. (2015). Successful aging 2.0: Conceptual expansions for the 21st century. *The Journals of Gerontology. Series B: Psychological Sciences and Social Sciences, 70*, 593–596. <http://dx.doi.org/10.1093/geronb/gbv025>

- Ryff, C. D., & Singer, B. (1998). The contours of positive human health. *Psychological Inquiry*, 9, 1–28. http://dx.doi.org/10.1207/s15327965pli0901_1
- Schilling, O. K., Wahl, H.-W., & Wiegering, S. (2013). Affective development in advanced old age: Analyses of terminal change in positive and negative affect. *Developmental Psychology*, 49, 1011–1020. <http://dx.doi.org/10.1037/a0028775>
- Siedler, T., Schupp, J., & Wagner, G. G. (2011). Innovative methods within in the context of secondary data: Examples from household panel surveys. In K. H. Trzesniewski, M. B. Donnellan, & R. E. Lucas (Eds.), *Secondary data analysis. An introduction to psychologists* (pp. 103–118). Washington, DC: American Psychological Association. <http://dx.doi.org/10.1037/12350-006>
- Silverstein, M., & Bengtson, V. L. (1994). Does intergenerational social support influence the psychological well-being of older parents? The contingencies of declining health and widowhood. *Social Science & Medicine*, 38, 943–957. [http://dx.doi.org/10.1016/0277-9536\(94\)90427-8](http://dx.doi.org/10.1016/0277-9536(94)90427-8)
- Singer, J. D., & Willett, J. B. (2003). *Applied longitudinal data analysis: Modeling change and event occurrence*. New York, NY: Oxford University Press. <http://dx.doi.org/10.1093/acprof:oso/9780195152968.001.0001>
- Sneed, R. S., & Cohen, S. (2013). A prospective study of volunteerism and hypertension risk in older adults. *Psychology and Aging*, 28, 578–586. <http://dx.doi.org/10.1037/a0032718>
- Snijders, T. A. B., & Bosker, R. J. (1999). *Multilevel analysis: An introduction to basic and advanced multilevel modeling*. London, England: Sage.
- Taylor, S. E. (2011). Social support: A review. In H. S. Friedman (Ed.), *The Oxford handbook of health psychology* (pp. 189–214). New York, NY: Oxford University Press.
- Uchino, B. N. (2006). Social support and health: A review of physiological processes potentially underlying links to disease outcomes. *Journal of Behavioral Medicine*, 29, 377–387. <http://dx.doi.org/10.1007/s10865-006-9056-5>
- Vogel, N., Schilling, O. K., Wahl, H. W., Beekman, A. T., & Penninx, B. W. (2013). Time-to-death-related change in positive and negative affect among older adults approaching the end of life. *Psychology and Aging*, 28, 128–141. <http://dx.doi.org/10.1037/a0030471>
- Windsor, T. D., Gerstorf, D., & Luszcz, M. A. (2015). Social resource correlates of levels and time-to-death-related changes in late-life affect. *Psychology and Aging*, 30, 136–148. <http://dx.doi.org/10.1037/a0038757>
- Zarit, S. H., & Reamy, A. M. (2013). Future directions in family and professional caregiving for the elderly. *Gerontology*, 59, 152–158. <http://dx.doi.org/10.1159/000342242>
- Zautra, A. J., Infurna, F. J., Zautra, E., Gallardo, C., & Velasco, L. (2016). The humanization of social relations: Nourishment for resilience. In A. Ong & C. E. Löckenhoff (Eds.), *Emotion, aging, and health*. Washington, DC: American Psychological Association.

Received July 7, 2015

Revision received November 25, 2015

Accepted December 7, 2015 ■